



# A contemporary inventory of gathered wild food plants in Liguria and Piedmont (NW Italy) and its implications for the in situ conservation of plant genetic resources

Ajmal Khan Manduzai · Andrea Pieroni

Received: 1 August 2026 / Accepted: 7 September 2026  
© The Author(s), under exclusive licence to Springer Nature B.V. 2026

**Abstract** National inventories of crop wild relatives (CWR) record where a wild relative occurs, but not whether anyone still gathers, manages, or discriminates it. Local knowledge of wild food plants supplies that missing attribute. We pooled four ethnobotanical datasets collected between 2024 and 2026 in three areas of Piedmont, the Chierese, the province of Cuneo, and the Monferrato, and in the Genoese area of Liguria, where the mixed wild-greens preparation known as *Preboggion* is still sold in the city market. Semi-structured interviews and field walks with 48 informants aged 24–89, complemented by participant observation and market surveys, yielded 62 wild taxa in 30 families. Forty-three (69%) fall within the taxon group definition of a CWR, and 28 are conspecific with a cultivated plant, including *Beta vulgaris* subsp. *maritima* (L.) Arcang., progenitor of all cultivated beets and the identity core of *Preboggion*, *Humulus lupulus* L., *Corylus avellana* L. and *Prunus avium* (L.) L.. Every taxon whose use is expanding is

a CWR, and no wild-harvested plant without a cultivated counterpart is gaining ground. Only *Urtica dioica* L. and *Taraxacum officinale* F.H.Wigg. were recorded in all four areas, and just under 70% of taxa in one only. In-field transmission has collapsed almost everywhere, and we documented three cases of incipient domestication. We argue that the erosion of foraging knowledge carries a genetic resources cost distinct from its cultural cost, and propose that ethnobotanical surveys become a routine input to CWR inventories.

**Keywords** Crop wild relatives · Wild food plants · Local ecological knowledge · Ethnobotany

## Introduction

Wild relatives of crops are the largest accessible reservoir of adaptive variation available to plant breeding, and their systematic conservation has been a stated international priority for two decades (Maxted et al. 2006, 2012; Heywood et al. 2007). The instruments developed to pursue that priority are, almost without exception, taxonomic and spatial. A national inventory of crop wild relatives (CWR) is assembled by applying the gene pool or taxon group concept to a national checklist, filtering by crop importance and threat status, and then mapping the resulting taxa (Vincent et al. 2013). Italy has such an inventory: Landucci et al. (2014) produced a full CWR and

---

A. K. Manduzai (✉)  
Research Department, Mili University, Airport 80 M Road,  
Khair Khana, Lab-E-Jar Square, Kabul, Afghanistan  
e-mail: ajmalmanduzai@gmail.com

A. Pieroni  
University of Gastronomic Sciences, Piazza Vittorio  
Emanuele II 9, 12042 Pollenzo, Italy

A. Pieroni  
Department of Medical Analysis, Tishk International  
University, Erbil 4001, Iraq

wild-harvested plant checklist for the Italian vascular flora, prioritised it, and subsequent literature has refined the priority list and assessed the conservation status of its members (Perrino and Perrino 2020). Gap analyses built on this logic have shown, at a global scale, how thinly the resulting diversity is represented in genebanks (Castañeda-Álvarez et al. 2016), and how consequential the material already used turned out to be (Dempewolf et al. 2017).

This approach does not record whether anyone still uses the taxon. A distribution polygon states that a wild relative occurs somewhere; it does not state that a community walks that slope every March, that it distinguishes two morphotypes within the stand and prefers one of them, that the harvest supplies a named product with an urban market, or that the habitat exists at all only because of a grazing regime that is now being abandoned. Those are ethnobotanical facts, and they bear directly on in situ conservation, because a genetic reserve is a management decision about people as much as about plants (Heywood et al. 2007). The point has been made in general terms that wild foods are actively managed, and the dichotomy between the cultivated and the wild is largely an artefact of how statistics are collected (Bharucha and Pretty 2010), but it is rarely operationalised. CWR checklists and wild food plant inventories are still, in practice, compiled by different communities of researchers who cite each other sparingly.

There is a second reason to bring the two literatures together. Domestication is not a completed historical event but an ongoing process, and wild food plants are its raw material (von Wettberg et al. 2020). The triage that identifies plausible candidates for neo-domestication—cultural salience, existing market value, agronomic tractability—takes its first two criteria directly from ethnobotanical data (Sulaiman et al. 2023). Where a wild food plant is being transplanted into kitchen gardens, moved from field margin to seedbed, or brought into cultivation to supply the herbal-products industry, the early stages of domestication are visible in real time, and the record of them is held by the people doing it.

Both arguments run into the same obstacle. Across Europe, the knowledge in question is disappearing faster than it is being written down. Reviews of contemporary wild food plant use have documented the collapse of subsistence-driven traditions alongside the emergence of new, aesthetically driven cuisines

built on wild edibles (Łuczaj et al. 2012; Turner et al. 2011), and Italy in particular has been mapped in detail, with hundreds of taxa recorded in traditional vegetable mixtures (Ghirardini et al. 2007; Sansanelli and Tassoni 2014; Guarrera and Savo 2016). Diachronic investigations that revisit the same territory after decades consistently report contraction: in the Southern Occitan Alps the number of families in active use narrowed markedly between the 1970s and 2022 (Alrhoun et al. 2025b), in the upper Susa Valley half a century of change produced a similar picture (Sulaiman et al. 2024b), in the western Alps the medicinal repertoire of the pastoral economy has largely given way to synanthropic taxa (Alrhoun et al. 2025c), and in central Tuscany a twenty-year comparison recorded a clear reduction in recognised taxa (Alrhoun et al. 2025a). In three Mediterranean coastal areas, the erosion has gone so far that the authors describe their search for a living foraging tradition as a search for a ghost (Sulaiman et al. 2026).

North-West Italy is a good place to examine these processes together. Its foraging traditions are unusually well described for the Alpine margin (Pieroni and Giusti 2009; Cornara et al. 2009; Vitalini et al. 2013); two of its wild plant preparations, the Ligurian *Preboggion*, a variable mixture of wild greens, and the Piedmontese *Luvertin*, the young shoots of wild hop, *Humulus lupulus* L., have survived as named, marketable culinary artefacts rather than as loose knowledge about individual species. Within less than two hundred kilometres, the region contains high Occitan valleys at 2,000 m, an intensively cultivated plain, peri-urban hills under pressure from Turin, and a dense coastal city where wild greens are sold daily in a covered market. It is also a region of considerable agricultural specialisation, so that several of the wild taxa gathered there are the conspecific wild populations of locally important crops.

This paper draws together four independent field datasets collected in these settings between 2024 and 2026. Our aims are: (i) to compile an inventory of the wild taxa currently or, until recent times, traditionally gathered across the four areas; (ii) to assess how many of them are crop wild relatives or wild harvested plants, and what the ethnobotanical record adds to what a checklist already states about them; (iii) to describe how gathering practices and the transmission of gathering knowledge have changed over the last two to three generations, as reported by the

gatherers themselves; and (iv) to draw out the consequences of that change for the in situ conservation and future use of the genetic resources concerned.

## Materials and methods

### Study areas


Four areas were investigated (Fig. 1). The Chierese (CH) is the south-eastern portion of the Metropolitan City of Turin, a transition zone between the Po plain and the Turin hills, between roughly 200 and 600 m a.s.l., historically farmed by small family holdings and now increasingly peri-urban; cherry orchards, vineyards, arable fields, residual woodland and field margins form a fine-grained agro-ecological mosaic.

The province of Cuneo (CN) is the largest and most heterogeneous area, and was investigated at two contrasting altitudes: the Occitan-speaking valleys of Varaita, Maira and Po, where gathering

was historically tied to a pastoral economy and to the cross-border herb trade towards France, and the plain around Racconigi, Caramagna, Savigliano and Cavallermaggiore, where intensive, largely monocultural agriculture has replaced the older mosaic.

The Monferrato area (MO) was investigated around Cioccaro and Penango in the province of Asti, a hill landscape of vineyards, fields and woodland with an ageing resident population, and with a two-Michelin-star restaurant operating within the same municipality, an unusually direct juxtaposition of domestic and haute-cuisine uses of the same plant.

The Genoese area (GE) comprises the rural quarter of Bavari, on the hills above the city, where traditional horticulture persists, and the Mercato Orientale in central Genoa, one of the city's main covered markets, where *Preboggion* is sold ready-mixed by specialised vendors. This pairing allowed us to follow the same plants from the gathering ground to the urban point of sale.



**Fig. 1** Study areas. **a** The four areas investigated in Piedmont and Liguria, NW Italy. CH = Chierese (Metropolitan City of Turin); CN = province of Cuneo, investigated at two contrasting altitudes (the Occitan-speaking valleys of Varaita, Maira and Po, and the plain around Racconigi, Caramagna, Savigliano and Cavallermaggiore); MO = Monferrato (province of Asti, around Cioccaro and Penango); GE = Genoa and its hinterland (Liguria). Province boundaries are shown in grey and

provincial names in italics; the red rectangle marks the extent of panel **b**. **b** The Ligurian study area, within the municipality of Genoa: the Mercato Orientale in the city centre, where *Preboggion* is sold ready-mixed by specialised vendors, and the rural quarter of Bavari in the hinterland above the city, where it is gathered. **c** Position of Piedmont and Liguria within Italy, with the two regions highlighted in red

## Field data collection

Fieldwork was carried out between 2024 and 2026. Data were collected through semi-structured interviews, informal group conversations and focus groups, participant observation during gathering walks, and market surveys, following standard ethnobotanical practice. In total 48 people were interviewed: 11 in the Genoese area (5 male, 6 female, aged 65–82), 20 in the province of Cuneo (5 male, 15 female; traditional knowledge holders aged 65–81 and restaurateurs aged 30–50), 6 in the Monferrato (3 male, 3 female, aged 38–81), and 11 in the Chierese (6 male, 5 female, aged 24–89). We purposively selected informants for their direct, long-term experience of gathering. They were identified through the field researchers' local knowledge, each of whom worked in a familiar area, and through referrals from earlier participants; in Genoa, the vendors of the Mercato Orientale were approached directly at their stalls. They included farmers, market gardeners, nursery growers, retired agricultural workers, herb sellers, a biologist specialised in *Preboggion*, an agronomist, and chefs. The purposive design aims at depth rather than at representativeness, and three measures were taken to limit the biases it entails: the four datasets were collected independently by different researchers; informant accounts were triangulated with participant observation during gathering walks and with market surveys; and the sample deliberately spans both genders, a wide age range and lay as well as professional knowledge holders. The residual limitations of this sampling are discussed in Sect. 4.8. Overall, 29 of the 48 informants (60%) were female, a predominance consistent with the largely female and domestic channel through which gathering knowledge was traditionally transmitted in the region (Sects. 3.5 and 4.6). Interviews were conducted in Italian, in Piedmontese and Genoese dialect, and in Occitan where appropriate. Local plant names were elicited in the informant's own language and, wherever possible, in the presence of the plant itself, during gathering walks or on fresh material at the market; the procedure by which vernacular names were linked to scientific taxa is described in Sect. 2.3.

Interview schedules differed slightly between areas but converged on the same core domains: local wild plant names; gathering sites and seasons; parts used; culinary, medicinal (internally administered) and

other domestic preparations; the boundary between “resource” and “weed”; whether the plant was ever transplanted, sown, tended or otherwise managed; how gathering was learnt and whether it is being passed on; and perceived changes in plant availability. Where informants agreed, we audio-recorded and transcribed interviews; otherwise, we took notes during and immediately after the encounter. Free listing of the components of *Preboggion* was used in the Genoese area, and citation frequencies were calculated on that sample. The consolidated interview guide, covering the core domains addressed in all four areas, is provided as Online Resource 1.

## Plant identification, voucher specimens and nomenclature

We identified plants in the field using morphological characters and habitat, with photographic documentation. Preliminary field determinations were supported by the PI@ntNet application (Joly et al. 2016) and were then verified against Flora d'Italia (Pignatti et al. 2017–2019).

No new voucher specimens were collected during the present study, which was designed as a non-extractive survey of use and knowledge rather than a collecting mission. Field determinations were instead cross-checked against the voucher specimens collected and deposited by our research group during two earlier collecting expeditions in the same and immediately adjoining areas: the upper Varaita valley (Pieroni and Giusti 2009) and the Brigasc, Kye and Provencal communities of the Gesso, Grana and Ellero valleys (Mattalia et al. 2013). Those specimens are deposited at the Herbarium of the University of Gastronomic Sciences, Pollenzo, Italy, and the voucher code of each taxon is reported in the species table of the corresponding publication; the codes are given in Table 1. We used three further diachronic investigations carried out by our group in the same portion of the Western Alps to compare uses and folk names. Still, they did not themselves generate vouchers: the upper Borbera valley (Fontefrancesco et al. 2022), where plants were identified from informants' own specimens, the upper Susa valley (Sulaiman et al. 2024b), which relied on material deposited during the earlier fieldwork, and the Southern Occitan valleys of Maira, Stura and Grana (Alrhoun et al. 2025b), which re-analysed existing datasets.

**Table 1** Wild taxa gathered in the four study areas of Liguria and Piedmont, NW Italy. Botanical nomenclature follows Plants of the World Online (POWO); identifications were made as described in the Methods. Area codes: CH=Chierese (Turin); CN=province of Cuneo; MO=Monferrato (Asti); GE=Genoa and Bavari (Liguria). Trend, as reported by informants: + increasing or newly commercialised; = persisting; – declining or no longer gathered. For taxa lacking a voucher specimen, identification rests on the convergent field evidence described in Sect. 2.3, and the record may also encompass other morphologically similar congenic species gathered under the same folk name

| Family         | Recorded taxon                                                                                   | Local folk name(s)                         | Part(s) used          | Recorded traditional and current uses                                                                 | Areas; trend | Voucher        |
|----------------|--------------------------------------------------------------------------------------------------|--------------------------------------------|-----------------------|-------------------------------------------------------------------------------------------------------|--------------|----------------|
| Asteraceae     | <i>Achillea erba-rotta</i> All. subsp. <i>moschata</i> (Wulfen) I.Richardson                     | <i>Canamio</i> , <i>Erba rotta</i> (CN)    | Flowering tops        | Aromatic herbal preparations (substitute for <i>genepti</i> ); evening tisane for sleep and digestion | CN =         | UNISG-CHI-CAN  |
| Amaryllidaceae | <i>Allium schoenoprasum</i> L                                                                    | <i>Erba cipollina selvatica</i> (CN)       | Leaves                | Flavouring; preserved in walnut oil                                                                   | CN –         | UNISG-CHI-POU  |
| Amaryllidaceae | <i>Allium ursinum</i> L                                                                          | <i>Aglia orsino</i> (CN, MO)               | Leaves                | Aioli-type sauce for boiled vegetables and cheese; preserved in oil; recent pesto-type preparations   | CN, MO +     | —              |
| Amaryllidaceae | <i>Allium vineale</i> L                                                                          | <i>Aj sarvaj</i> (CH)                      | Bulbs, leaves         | Flavouring                                                                                            | CH –         | —              |
| Rosaceae       | <i>Aria edulis</i> (Willd.) M.Roem                                                               | <i>Sorbo montano</i> (CN)                  | Fruits                | Betted under straw and then eaten                                                                     | CN –         | —              |
| Asteraceae     | <i>Artemisia absinthium</i> L                                                                    | <i>Assenzio</i> (CN)                       | Aerial parts          | Aromatic herbal preparations; tisanes (used empirically during the 1918 influenza pandemic)           | CN –         | UNISG-CHI-BIO  |
| Asteraceae     | <i>Artemisia genipi</i> Weber ex Stechm., <i>A. glacialis</i> L. and <i>A. umbelliformis</i> Lam | <i>Genepti maschi</i> , <i>Fumelo</i> (CN) | Flowering tops        | <i>Genepti</i> aromatic herbal preparation; formerly traded across the French border                  | CN =         | UNISG-CHI-GEF2 |
| Asteraceae     | <i>Artemisia vulgaris</i> L                                                                      | <i>Artemisia comune</i> (CN)               | Aerial parts          | Historical bulk herb trade                                                                            | CN –         | UNISG-CHI-ART  |
| Asparagaceae   | <i>Asparagus officinalis</i> L                                                                   | <i>Asparagi selvatici</i> (CH)             | Young shoots          | Boiled, omelettes; currently sold at farmers' markets                                                 | CH +         | —              |
| Asteraceae     | <i>Bellis perennis</i> L                                                                         | <i>Margheritina</i> (MO)                   | Whole plant with root | Minestra di fiori (spring soup of mixed wild greens and flowers)                                      | MO –         | —              |
| Amaranthaceae  | <i>Beta vulgaris</i> subsp. <i>maritima</i> (L.) Arcang                                          | <i>Gea</i> , <i>Bietoa sarvaega</i> (GE)   | Leaves                | Core ingredient of preboggion; soups, savoury pies, pansoti and ravioli fillings                      | GE =         | —              |
| Amaranthaceae  | <i>Blitum bonus-henricus</i> (L.) Rchb                                                           | <i>Spinaci selvatici</i> (CN)              | Leaves, shoots        | Boiled as a vegetable; fillings for capunet                                                           | CN –         | UNISG-CHI-BIE  |

Table 1 (continued)

| Family       | Recorded taxon                                       | Local folk name(s)                                                | Part(s) used               | Recorded traditional and current uses                                                                                                                              | Areas; trend | Voucher       |
|--------------|------------------------------------------------------|-------------------------------------------------------------------|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|---------------|
| Boraginaceae | <i>Borago officinalis</i> L                          | <i>Borragine</i> (GE, CN, MO)                                     | Leaves, flowers            | Preboggion; agnolotti and ravioli fillings; fritters; tisane for headache                                                                                          | GE, CN, MO = | —             |
| Cornaceae    | <i>Cornus mas</i> L                                  | <i>Comiolo</i> (CH)                                               | Fruits                     | Eaten raw; preserves                                                                                                                                               | CH –         | —             |
| Betulaceae   | <i>Corylus avellana</i> L                            | <i>Nocciolo</i> (CN)                                              | Seeds                      | Pressed for oil used to dress wild greens                                                                                                                          | CN –         | UNISG-DIS025  |
| Apiaceae     | <i>Foeniculum vulgare</i> Mill                       | <i>Finocchietto selvatico</i> (GE)                                | Shoots, leaves             | Preboggion; flavouring                                                                                                                                             | GE =         | —             |
| Rosaceae     | <i>Fragaria vesca</i> L                              | <i>Fragoline di bosco</i> (CH, CN, MO)                            | Fruits, leaves             | Eaten raw; jams; leaves in minestra di fiori; plants transplanted into kitchen gardens                                                                             | CH, CN, MO = | UNISG-CHI-MOU |
| Gentianaceae | <i>Gentiana acaulis</i> L                            | <i>Gensianella, Chause del cucù</i> (CN)                          | Flowers                    | Decoctions and tisanes, including for evening anxiety                                                                                                              | CN –         | UNISG-CHI-CHC |
| Gentianaceae | <i>Gentiana lutea</i> L                              | <i>Genziana maggiore</i> (CN)                                     | Roots                      | Bitter digestive preparations                                                                                                                                      | CN =         | UNISG-CHI-ERG |
| Cannabaceae  | <i>Humulus lupulus</i> L                             | <i>Luvertin, Uvertin, Invertin</i> (MO); <i>Luvertin</i> (CN, CH) | Young shoots; female cones | Shoots steamed, in omelettes, risotti, fresh-pasta fillings and salads; cones for flavouring beverages and for sedative infusions; local fair at Sant'Albano Stura | CH, CN, MO + | UNISG-DIS010  |
| Asteraceae   | <i>Hyoseris radiata</i> L                            | <i>Piscianletto, Garatun</i> (GE)                                 | Leaf rosettes              | Preboggion                                                                                                                                                         | GE =         | —             |
| Juglandaceae | <i>Juglans regia</i> L                               | <i>Noci</i> (CN)                                                  | Seeds                      | Pressed for oil used to dress wild greens; flavoured beverages                                                                                                     | CN –         | —             |
| Asteraceae   | <i>Leontodon hispidus</i> L                          | <i>Tageinetti</i> (GE)                                            | Leaf rosettes              | Preboggion                                                                                                                                                         | GE –         | —             |
| Asteraceae   | <i>Leontopodium nivale</i> (Ten.) Huet ex Hand.-Mazz | <i>Stella alpina</i> (CN)                                         | Flower heads               | Aromatic herbal preparations (species under legal protection; gathering now prohibited)                                                                            | CN –         | —             |
| Malvaceae    | <i>Malva sylvestris</i> L                            | <i>Malva</i> (GE, CH, CN)                                         | Leaves, flowers            | Preboggion; soups; emollient tisane                                                                                                                                | GE, CH, CN = | UNISG-CHI-RIU |
| Asteraceae   | <i>Matricaria chamomilla</i> L                       | <i>Camomilla</i> (CH, CN)                                         | Flower heads               | Tisanes; in the Raconigi plain shifted from field weed to transplanted crop                                                                                        | CH, CN +     | UNISG-DIS004  |

**Table 1** (continued)

| Family          | Recorded taxon                                         | Local folk name(s)                          | Part(s) used    | Recorded traditional and current uses                                                                              | Areas; trend | Voucher        |
|-----------------|--------------------------------------------------------|---------------------------------------------|-----------------|--------------------------------------------------------------------------------------------------------------------|--------------|----------------|
| Lamiaceae       | <i>Melissa officinalis</i> L                           | <i>Melissa</i> (CH, CN)                     | Leaves          | Tisanes; lemon note in dandelion-flower syrup; capunet fillings                                                    | CH, CN =     | —              |
| Lamiaceae       | <i>Mentha spicata</i> L                                | <i>Menta selvatica</i> (CN)                 | Leaves          | Omelettes, soups, syrup, flavoured beverages; frozen syrup ( <i>Toc de giasa</i> ); later cultivated at Pancalieri | CN =         | —              |
| Moraceae        | <i>Morus nigra</i> L                                   | <i>Moro, Gelso</i> (CH, MO)                 | Fruits          | Eaten raw; preserves                                                                                               | CH, MO –     | —              |
| Brassicaceae    | <i>Nasturtium officinale</i> W.T.Aiton                 | <i>Crescione delle fontane</i> (CN)         | Leaves          | Salads; flavouring for fresh cheeses                                                                               | CN –         | UNISG-CHI-CRE  |
| Papaveraceae    | <i>Papaver rhoeas</i> L                                | <i>Papavau</i> (GE); <i>Rosolacci</i> (CH)  | Leaf rosettes   | Preboggion; boiled greens; explicitly recalled as a hunger filler                                                  | GE, CH –     | —              |
| Urticaceae      | <i>Parietaria officinalis</i> L                        | <i>Erba vetriola</i> (CH)                   | Leaves          | Occasional food; minor medicinal use                                                                               | CH –         | —              |
| Pinaceae        | <i>Pinus cembra</i> L                                  | <i>Pinoli, Pino cembro</i> (CN)             | Seeds           | Pressed for oil; source of unsaturated fats at high altitude                                                       | CN –         | UNISG-CHI-ELV  |
| Plantaginaceae  | <i>Plantago lanceolata</i> L                           | <i>Piantaggine</i> (GE, CH)                 | Leaves          | Preboggion; minor medicinal uses                                                                                   | GE, CH –     | UNISG-CHI-PIA1 |
| Plantaginaceae  | <i>Plantago major</i> L                                | <i>Piantaggine rotonda</i> (GE)             | Leaves          | Preboggion                                                                                                         | GE –         | UNISG-CHI-PIA2 |
| Portulacaceae   | <i>Portulaca oleracea</i> L                            | <i>Erba porcellana</i> (CH)                 | Shoots, leaves  | Salads; currently sold at farmers' markets                                                                         | CH +         | —              |
| Primulaceae     | <i>Primula veris</i> L. (and <i>P. vulgaris</i> Huds.) | <i>Blins, Primule</i> (CN, MO)              | Leaves, flowers | Omelettes on feast days; minestra di fiori                                                                         | CN, MO –     | UNISG-DIS032   |
| Rosaceae        | <i>Prunus avium</i> (L.) L                             | <i>Ciliegie selvatiche</i> (MO)             | Fruits          | Eaten raw; preserves                                                                                               | MO =         | UNISG-DIS034   |
| Asteraceae      | <i>Reichardia picroides</i> (L.) Roth                  | <i>Talegua, Gattalevre</i> (GE)             | Leaf rosettes   | Preboggion; raw in salads                                                                                          | GE =         | —              |
| Grossulariaceae | <i>Ribes rubrum</i> L                                  | <i>Ribes</i> (CN)                           | Fruits          | Jams; seedlings transplanted into kitchen gardens                                                                  | CN –         | —              |
| Fabaceae        | <i>Robinia pseudoacacia</i> L                          | <i>Fiur de ghersia, Fiori d'acacia</i> (CN) | Flowers         | Fritters served with honey                                                                                         | CN =         | UNISG-DIS019   |
| Rosaceae        | <i>Rosa canina</i> L                                   | <i>Gratacul</i> (CN)                        | Hips            | Jams, after removal of seeds and hairs                                                                             | CN –         | UNISG-CHI-BUC  |

Table 1 (continued)

| Family          | Recorded taxon                                          | Local folk name(s)                                                                 | Part(s) used             | Recorded traditional and current uses                                                                                                   | Areas; trend     | Voucher        |
|-----------------|---------------------------------------------------------|------------------------------------------------------------------------------------|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|------------------|----------------|
| Rosaceae        | <i>Rubus ulmifolius</i> Schott                          | <i>More, Rovo</i> (CN, MO)                                                         | Fruits, young shoots     | Jams; a traditional beverage ( <i>Vinus moretus</i> ); shoots in minestra di fiori                                                      | CN, MO =         | —              |
| Polygonaceae    | <i>Rumex acetosa</i> L.                                 | <i>Acetosà</i> (CH)                                                                | Leaves                   | Eaten raw as a sour snack; flavoured                                                                                                    | CH –             | UNISG-CHI-SET1 |
| Polygonaceae    | <i>Rumex alpinus</i> L.                                 | <i>Rabarbaro selvatico</i> (CN)                                                    | Leaves, petioles         | Leaf wrappers for capunet when cabbage was unavailable                                                                                  | CN –             | UNISG-CHI-GRA  |
| Lamiaceae       | <i>Salvia pratensis</i> L.                              | <i>Salvia dei prati</i> (CH)                                                       | Leaves                   | Flavoured                                                                                                                               | CH –             | UNISG-DIS023   |
| Viburnaceae     | <i>Sambucus nigra</i> L.                                | <i>Sambuco</i> (CH, CN)                                                            | Flowers, fruits          | Fritters, flavoured beverages, tisanes; berry juice for syrup, jam and cordial                                                          | CH, CN =         | UNISG-DIS012   |
| Rosaceae        | <i>Sanguisorba minor</i> Scop                           | <i>Pimpinella</i> (GE)                                                             | Leaves                   | Preboggion; raw in salads                                                                                                               | GE =             | —              |
| Caryophyllaceae | <i>Silene vulgaris</i> (Moench) Gareke                  | <i>Scipuelli</i> (GE); <i>Coujet, Cuiet</i> (CH, CN)                               | Young shoots, leaves     | Preboggion; risotti, omelettes, minestra d'erbette                                                                                      | GE, CH, CN =     | UNISG-DIS011   |
| Brassicaceae    | <i>Sinapis arvensis</i> L.                              | <i>Senape selvatica</i> (GE, MO)                                                   | Leaves                   | Preboggion; boiled greens                                                                                                               | GE, MO –         | —              |
| Asteraceae      | <i>Sonchus oleraceus</i> L.                             | <i>Scixerboa, Cardelle</i> (GE)                                                    | Leaf rosettes            | Preboggion                                                                                                                              | GE =             | —              |
| Asteraceae      | <i>Taraxacum officinale</i> F.H.Wigg                    | <i>Latussa</i> (GE); <i>Muriun, Scioria di pra</i> (CN); <i>Tarassaco</i> (CH, MO) | Rosettes, roots, flowers | Raw in salads when young, boiled when mature; depurative decoction; roasted root as coffee surrogate; flower syrup (amel de virasuleil) | GE, CH, CN, MO = | UNISG-CHI-MOU  |
| Lamiaceae       | <i>Thymus serpyllum</i> L. s.l                          | <i>Serpoul, Timo</i> (CN)                                                          | Aerial parts             | Flavoured meat and Saras cheese; tisanes; used to mask the taste of marmot meat                                                         | CN =             | UNISG-DIS024   |
| Asteraceae      | <i>Tragopogon pratensis</i> L.                          | <i>Barbabuch, Barbaboc</i> (CH, MO)                                                | Roots, young shoots      | Boiled or in salads; a wartime and post-war staple                                                                                      | CH, MO –         | UNISG-CHI-BAR  |
| Fabaceae        | <i>Trifolium pratense</i> L.                            | <i>Trifoglio dei prati</i> (CH)                                                    | Flowers, leaves          | Occasional food; fodder                                                                                                                 | CH –             | —              |
| Asteraceae      | <i>Urospermum dalechampii</i> (L.) Scop. ex F.W.Schmidt | <i>Bell'omu</i> (GE)                                                               | Leaf rosettes            | Preboggion                                                                                                                              | GE =             | —              |
| Urticaceae      | <i>Urtica dioica</i> L.                                 | <i>Ortica</i> (GE, CH, CN, MO)                                                     | Young shoots             | Omelettes, soups, fillings; preboggion; gathered by children in spring                                                                  | GE, CH, CN, MO = | UNISG-CHI-URT  |

**Table 1** (continued)

| Family         | Recorded taxon                                                            | Local folk name(s)                               | Part(s) used  | Recorded traditional and current uses                                                                   | Areas; trend | Voucher       |
|----------------|---------------------------------------------------------------------------|--------------------------------------------------|---------------|---------------------------------------------------------------------------------------------------------|--------------|---------------|
| Ericaceae      | <i>Vaccinium myrtillus</i> L.                                             | <i>Asaires</i> , <i>Mirtilli neri</i> (CN)       | Fruits        | Raw and cooked; preserves; flavoured beverages                                                          | CN =         | UNISG-DIS017  |
| Ericaceae      | <i>Vaccinium vitis-idaea</i> L.                                           | <i>Mirtilli rossi</i> (CN)                       | Fruits        | Historically rejected as indigestible; eaten only after cooking                                         | CN –         | —             |
| Caprifoliaceae | <i>Valerianella locusta</i> (L.) Laterr                                   | <i>Salezet</i> , <i>Valeriana selvatica</i> (CN) | Leaf rosettes | Salads dressed with walnut or stone-pine oil                                                            | CN –         | —             |
| Plantaginaceae | <i>Veronica cf. allionii</i> Vill                                         | <i>Tè di montagna</i> (CN)                       | Aerial parts  | High-altitude substitute for tea                                                                        | CN –         | UNISG-CHI-IAS |
| Violaceae      | <i>Viola odorata</i> L. (and <i>V. calcarata</i> L. in the upper valleys) | <i>Violette</i> (CN, MO)                         | Flowers       | Decoctions; violet cream and syrup; formerly gathered by girls and sold to confectioners and herbalists | CN, MO –     | UNISG-CHI-VIO |

For the taxa in Table 1 for which no voucher specimen exists, identification rests on three converging lines of field evidence. The first is photographic documentation of the plant in situ, taken at the phenological stage at which it is actually gathered, since several of the taxa concerned are collected at rosette or young-shoot stage and are hardest to determine precisely then. The second is the morphological and ecological description given by the informant, covering habitat, slope, season and, in particular, the characters the gatherers themselves use to discriminate the plant from its look-alikes. The third is the dialect name, compared both with the phytonyms elicited for the same taxa during our earlier expeditions in the same and adjoining areas (Pieroni and Giusti 2009; Mattalia et al. 2013) and with those documented in the ethnobotanical literature for Liguria, Piedmont and the Occitan Alps (Cornara et al. 2009; Fontefrancesco et al. 2022; Alrhoun et al. 2025b). Where these three lines did not converge unambiguously, the record was retained only at the level at which it was secure, that is at genus level or as a folk category spanning more than one botanical species, and is reported as such in Table 1. We regard this procedure as weaker evidence than a deposited specimen, and the distinction is visible in Table 1, where determinations resting on it can be told apart from those backed by a voucher.

We took particular care with taxa that co-occur in the study areas with morphologically similar species that non-specialists may find difficult to separate. The clearest case is the guild of edible-leaved Asteraceae gathered at rosette stage for *Preboggion* (*Hyoseris radiata* L., *Reichardia picroides* (L.) Roth, *Urospermum dalechampii* (L.) Scop. ex F.W.Schmidt, *Leontodon hispidus* L., *Sonchus oleraceus* L. and *Taraxacum officinale* F.H.Wigg.), which overlaps with several congeners and with other yellow-flowered composites of the same habitats; here identification relied precisely on the discriminating characters that the gatherers themselves use—bitterness, leaf texture and rosette shape—cross-checked against the photographs and the dialect names. *Allium ursinum* L., whose leaves are notoriously confused elsewhere in Europe with those of the toxic *Colchicum autumnale* L. and *Convallaria majalis* L., was always verified by smell and by the presence of flowering individuals in the stand. Where a folk category demonstrably spans more than one botanical species—as with the



**Fig. 2** *Artemisia genipi* Weber ex Stechm., one of the three species gathered under the folk category *Genepi*, on high-altitude scree in the Alpine belt of the province of Cuneo (Photo: A. Pieroni)



**Fig. 3** Flowering tops of *Thymus serpyllum* L. s.l. as gathered by informants in the province of Cuneo (Photo: A. Pieroni)

*Primula* and *Viola* complexes, with the folk category *Genepi*—which spans *Artemisia genipi* Weber ex Stechm., *A. glacialis* L. and *A. umbelliformis* Lam. (Fig. 2)—and with *Thymus serpyllum* s.l. (Fig. 3)—the record is reported at that broader level in Table 1 rather than forced onto a single binomial. For the taxa lacking a voucher specimen, we accordingly regard the determinations as referable to the species indicated, while acknowledging that other closely similar congeneric species may occasionally be involved; this is stated explicitly in the caption of Table 1.

Botanical nomenclature only was standardised afterwards: because the four datasets used different

naming conventions, each name was checked individually against Plants of the World Online (POWO; Govaerts et al. 2021) and, where the name used in the field was a synonym, replaced by the currently accepted one. POWO served as the nomenclatural authority and played no part in identifying the specimens. Author citations are given at the first mention of each binomial in the abstract and in the main text, and for every taxon in Tables 1 and 2. Families follow the same source. Several folk categories are deliberately broader than botanical species.

#### Assignment of CWR and wild harvested plant status

We assigned each of the 62 taxa in the inventory a CWR status following the taxon group concept of Maxted et al. (2006), which uses the existing taxonomic hierarchy as a proxy for relatedness where crossing and genetic diversity data are unavailable. Three classes were used. Taxa conspecific with a cultivated plant, that is, wild or feral populations of a species that is itself grown as a crop, were assigned to taxon group 1b (TG1b); taxa congeneric with a crop but not conspecific with it were assigned to taxon groups 2 to 4 (TG2–4), without attempting to resolve the section- and series-level distinctions, which are not available for all of the genera concerned; taxa with no cultivated congener were classed as wild harvested plants (WHP) in the sense of Landucci et al. (2014), that is, wild taxa of direct socio-economic value that are not relatives of a crop.

We identified the associated crop or crops from recorded field use, together with standard literature on Italian plant genetic resources (Landucci et al. 2014; Perrino and Perrino 2020). Two conventions should be stated. First, “crop” is used broadly, and includes food, aromatic, medicinal and beverage crops, since several of the taxa in our material are grown commercially in Piedmont for aromatic and herbal preparations rather than as vegetables; purely ornamental cultivation was not treated as sufficient. Second, the assignment is indicative rather than definitive: it rests on the taxon group concept and not on documented crossing behaviour, and for several genera the relationship with the crop gene pool is remote. Where a taxon is conspecific with a crop but the gathered populations are more plausibly feral escapes than genuinely wild ones, this is stated in Table 2 rather than silently resolved.

**Table 2** Crop wild relatives (CWR) and wild harvested plant status of the 62 taxa recorded, assigned following the taxon group concept of Maxted et al. (2006) as described in Sect. 2.4. TG1b=conspicuous with a cultivated plant, i.e. wild or feral populations of a crop species; TG2-4=congeneric with a crop but not conspecific with it; WHP=wild harvested plant with no cultivated congener. Area codes and trend symbols as in Table 1. Red List=category at the global and/or European level (Bilz et al. 2011; Allen et al. 2014; IUCN 2026), followed by the category in the national Red List of the Italian flora (Orsenigo et al. 2021): LC=Least Concern, NT=Near Threatened, NE=not evaluated, —=not assessed at the national level. No recorded taxon is threatened at any of these levels; the only national listing above LC is *Gentiana lutea* (NT). Regional harvest restrictions are indicated in the Notes column (Regione Piemonte 1982; Regione Liguria 2009)

| Taxon                                                                                            | Family         | Class | Related crop(s)                              | Areas; trend | Notes                                                                                                                                                       | Red List                                                                                       |
|--------------------------------------------------------------------------------------------------|----------------|-------|----------------------------------------------|--------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|
| <i>Achillea erba-rotta</i> All.<br>subsp. <i>moschata</i> (Wulfen)<br>I.Richardson               | Asteraceae     | TG1b  | Alpine aromatic herb (cultivated)            | CN =         | Brought into cultivation in Alpine areas for aromatic herbal preparations                                                                                   | NE / —                                                                                         |
| <i>Allium schoenoprasum</i> L.                                                                   | Amaryllidaceae | TG1b  | Chive                                        | CN –         | Wild populations of a species grown commercially as a herb                                                                                                  | LC / —                                                                                         |
| <i>Allium ursinum</i> L.                                                                         | Amaryllidaceae | TG2-4 | Onion, garlic, leek, chive                   | CN, MO +     | Increasingly cultivated in Europe as a leaf crop                                                                                                            | LC / —                                                                                         |
| <i>Allium vineale</i> L.                                                                         | Amaryllidaceae | TG2-4 | Onion, garlic, leek                          | CH –         | Also a weed of arable land                                                                                                                                  | LC / —                                                                                         |
| <i>Aria edulis</i> (Willd.) M.Roem                                                               | Rosaceae       | WHP   | —                                            | CN –         | Fruits bletted under straw                                                                                                                                  | LC / —                                                                                         |
| <i>Artemisia absinthium</i> L.                                                                   | Asteraceae     | TG1b  | Wormwood (aromatic and bitter preparations)  | CN –         | Cultivated commercially for the beverage industry                                                                                                           | LC / —                                                                                         |
| <i>Artemisia genipi</i> Weber ex Stechm., <i>A. glacialis</i> L. and <i>A. umbelliformis</i> Lam | Asteraceae     | TG1b  | <i>Genipi</i> aromatic herb                  | CN =         | Cultivation established in Alpine areas from wild germplasm; wild gathering subject to the authorisation regime for official plants (Regione Piemonte 1982) | <i>A. genipi</i> : LC / LC<br><i>A. glacialis</i> : NE / —<br><i>A. umbelliformis</i> : LC / — |
| <i>Artemisia vulgaris</i> L.                                                                     | Asteraceae     | TG2-4 | Wormwood, tarragon                           | CN –         | Historical bulk herb trade                                                                                                                                  | LC / —                                                                                         |
| <i>Asparagus officinalis</i> L.                                                                  | Asparagaceae   | TG1b  | Asparagus                                    | CH +         | Wild populations of the crop species; now sold at farmers' markets                                                                                          | LC / —                                                                                         |
| <i>Bellis perennis</i> L.                                                                        | Asteraceae     | WHP   | —                                            | MO –         | Meadow guild                                                                                                                                                | NE / —                                                                                         |
| <i>Beta vulgaris</i> subsp. <i>maritima</i> (L.) Arcang                                          | Amaranthaceae  | TG1b  | Beetroot, chard, sugar beet, fodder beet     | GE =         | Wild progenitor of all cultivated beets; identity core of Preboggion                                                                                        | LC / —                                                                                         |
| <i>Blitum bonus-henricus</i> (L.) Rchb                                                           | Amaranthaceae  | TG2-4 | Quinoa and allied Chenopodiaceae grain crops | CN –         | Remote relationship; nitrophilous stands tied to summer pastoral settlement                                                                                 | NE / —                                                                                         |
| <i>Borago officinalis</i> L.                                                                     | Boraginaceae   | TG1b  | Borage                                       | GE, CN, MO = | Cultivated in Liguria for pasta fillings; local landraces                                                                                                   | NE / —                                                                                         |
| <i>Cornus mas</i> L.                                                                             | Cornaceae      | TG1b  | Cornelian cherry                             | CH –         | Fruit cultivars selected in central and eastern Europe                                                                                                      | LC / —                                                                                         |

Table 2 (continued)

| Taxon                                                | Family       | Class | Related crop(s)                                  | Areas; trend | Notes                                                                                                                                                                                                              | Red List |
|------------------------------------------------------|--------------|-------|--------------------------------------------------|--------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|
| <i>Corylus avellana</i> L                            | Betulaceae   | TG1b  | Hazelnut                                         | CN –         | Wild populations of the crop underpinning the Piedmontese hazelnut sector                                                                                                                                          | LC / —   |
| <i>Foeniculum vulgare</i> Mill                       | Apiaceae     | TG1b  | Florence fennel, fennel seed                     | GE =         | Wild populations of the progenitor of cultivated fennel                                                                                                                                                            | LC / —   |
| <i>Fragaria vesca</i> L                              | Rosaceae     | TG2–4 | Garden strawberry ( <i>Fragaria x ananassa</i> ) | CH, CN, MO = | Also cultivated in its own right; plants transplanted into kitchen gardens                                                                                                                                         | LC / —   |
| <i>Gentiana acaulis</i> L                            | Gentianaceae | TG2–4 | Yellow gentian (bitter digestive preparations)   | CN –         | Congeneric with a cultivated bitter crop                                                                                                                                                                           | LC / —   |
| <i>Gentiana lutea</i> L                              | Gentianaceae | TG1b  | Yellow gentian (bitter digestive preparations)   | CN =         | Now cultivated; wild populations are the germplasm base; root collection regulated (Regione Piemonte 1982)                                                                                                         | LC / NT  |
| <i>Humulus lupulus</i> L                             | Cannabaceae  | TG1b  | Hop                                              | CH, CN, MO + | Wild populations of the crop species; the Luverrin of the inventory                                                                                                                                                | LC / —   |
| <i>Hyoseris radiata</i> L                            | Asteraceae   | WHP   | —                                                | GE =         | Preboggion component; two folk names in the Genoese sample                                                                                                                                                         | NE / —   |
| <i>Juglans regia</i> L                               | Juglandaceae | TG1b  | Walnut                                           | CN –         | Feral and planted populations; oil pressed for dressing wild greens                                                                                                                                                | LC / —   |
| <i>Leontodon hispidus</i> L                          | Asteraceae   | WHP   | —                                                | GE –         | Meadow guild                                                                                                                                                                                                       | NE / —   |
| <i>Leontopodium nivale</i> (Ten.) Huet ex Hand.-Mazz | Asteraceae   | WHP   | —                                                | CN –         | Historical use only; not on the regional absolute-protection list; gathering falls under the general five-specimen daily limit, and the law itself promotes the cultivation of the species (Regione Piemonte 1982) | LC / LC  |
| <i>Malva sylvestris</i> L                            | Malvaceae    | WHP   | —                                                | GE, CH, CN = | Preboggion component                                                                                                                                                                                               | LC / —   |
| <i>Matricaria chamomilla</i> L                       | Asteraceae   | TG1b  | Chamomile                                        | CH, CN +     | Documented shift from field weed to transplanted crop in the Raccomigi plain                                                                                                                                       | LC / —   |

**Table 2** (continued)

| Taxon                                                  | Family          | Class | Related crop(s)                           | Areas; trend | Notes                                                                            | Red List |
|--------------------------------------------------------|-----------------|-------|-------------------------------------------|--------------|----------------------------------------------------------------------------------|----------|
| <i>Melissa officinalis</i> L                           | Lamiaceae       | TG1b  | Lemon balm                                | CH, CN =     | Wild populations of a cultivated aromatic                                        | LC / —   |
| <i>Mentha spicata</i> L                                | Lamiaceae       | TG1b  | Spearmint, peppermint                     | CN =         | Later brought into cultivation at Pancalieri, a designated Italian mint district | LC / —   |
| <i>Morus nigra</i> L                                   | Moraceae        | TG1b  | Black mulberry                            | CH, MO –     | Feral and relict planted trees                                                   | NE / —   |
| <i>Nasturtium officinale</i> W.T.Aiton                 | Brassicaceae    | TG1b  | Watercress                                | CN –         | Wild populations of the crop species                                             | LC / —   |
| <i>Papaver rhoeas</i> L                                | Papaveraceae    | TG2–4 | Opium poppy ( <i>Papaver somniferum</i> ) | GE, CH –     | Arable guild; explicitly recalled as a hunger filler                             | NE / —   |
| <i>Parietaria officinalis</i> L                        | Urticaceae      | WHP   | —                                         | CH –         | Occasional food                                                                  | NE / —   |
| <i>Pinus cembra</i> L                                  | Pinaceae        | WHP   | —                                         | CN –         | Seed pressing for oil ceased with the trade                                      | LC / —   |
| <i>Plantago lanceolata</i> L                           | Plantaginaceae  | WHP   | —                                         | GE, CH –     | Meadow guild                                                                     | LC / —   |
| <i>Plantago major</i> L                                | Plantaginaceae  | WHP   | —                                         | GE –         | Preboggion component                                                             | LC / —   |
| <i>Portulaca oleracea</i> L                            | Portulacaceae   | TG1b  | Purslane                                  | CH +         | Wild populations of the crop; currently sold at farmers' markets                 | NE / —   |
| <i>Primula veris</i> L. (and <i>P. vulgaris</i> Huds.) | Primulaceae     | WHP   | —                                         | CN, MO –     | Montane meadow margin guild                                                      | LC / —   |
| <i>Prunus avium</i> (L.) L                             | Rosaceae        | TG1b  | Sweet cherry                              | MO =         | Wild populations of the crop; the Chierese is a cherry-orchard district          | LC / —   |
| <i>Reichardia picroides</i> (L.) Roth                  | Asteraceae      | WHP   | —                                         | GE =         | Candidate for neo-domestication as a leafy vegetable                             | NE / —   |
| <i>Ribes rubrum</i> L                                  | Grossulariaceae | TG1b  | Redcurrant                                | CN –         | Seedlings transplanted into kitchen gardens by informants                        | LC / —   |
| <i>Robinia pseudoacacia</i> L                          | Fabaceae        | WHP   | —                                         | CN =         | Naturalised alien; flowers used for fritters                                     | LC / —   |
| <i>Rosa canina</i> L                                   | Rosaceae        | TG2–4 | Cultivated roses (rootstock), rosehip     | CN –         | Standard rootstock for cultivated roses                                          | LC / —   |
| <i>Rubus ulmifolius</i> Schott                         | Rosaceae        | TG2–4 | Cultivated blackberry                     | CN, MO =     | Congeneric with, and interfertile within, the cultivated blackberry complex      | NE / —   |
| <i>Rumex acetosa</i> L                                 | Polygonaceae    | TG1b  | Garden sorrel                             | CH –         | Wild populations of a cultivated leaf vegetable                                  | LC / —   |

Table 2 (continued)

| Taxon                                                   | Family          | Class | Related crop(s)                           | Areas; trend     | Notes                                                                                                               | Red List |
|---------------------------------------------------------|-----------------|-------|-------------------------------------------|------------------|---------------------------------------------------------------------------------------------------------------------|----------|
| <i>Rumex alpinus</i> L                                  | Polygonaceae    | TG2–4 | Garden sorrel                             | CN –             | Nitrophilous stands dependent on transhumant husbandry                                                              | NE / —   |
| <i>Salvia pratensis</i> L                               | Lamiaceae       | TG2–4 | Sage ( <i>Salvia officinalis</i> )        | CH –             | Meadow guild; congeneric with a cultivated aromatic                                                                 | NE / —   |
| <i>Sambucus nigra</i> L                                 | Viburnaceae     | TG1b  | Elder (fruit and flower cultivars)        | CH, CN =         | Cultivars selected in central Europe                                                                                | LC / —   |
| <i>Sanguisorba minor</i> Scop                           | Rosaceae        | TG1b  | Salad burnet                              | GE =             | Cultivated forms exist; Preboggion component                                                                        | NE / —   |
| <i>Silene vulgaris</i> (Moench) Garcke                  | Caryophyllaceae | WHP   | —                                         | GE, CH, CN =     | Semi-cultivated elsewhere in southern Europe; neo-domestication candidate                                           | NE / —   |
| <i>Sinapis arvensis</i> L                               | Brassicaceae    | TG2–4 | Brassica oilseeds and vegetables, mustard | GE, MO –         | Arable margin guild; tertiary gene pool relationship                                                                | LC / —   |
| <i>Sonchus oleraceus</i> L                              | Asteraceae      | WHP   | —                                         | GE =             | Preboggion component; two folk names                                                                                | NE / —   |
| <i>Taraxacum officinale</i> F.H.Wigg                    | Asteraceae      | TG1b  | Dandelion (leaf and root cultivars)       | GE, CH, CN, MO = | Cultivated selections exist; recorded in all four study areas (along with <i>Urtica dioica</i> L.)                  | NE / —   |
| <i>Thymus serpyllum</i> L. s.l                          | Lamiaceae       | TG2–4 | Thyme ( <i>Thymus vulgaris</i> )          | CN =             | Congeneric with a cultivated aromatic                                                                               | NE / —   |
| <i>Tragopogon pratensis</i> L                           | Asteraceae      | TG2–4 | Salsify ( <i>Tragopogon porrifolius</i> ) | CH, MO –         | Congeneric with a root vegetable                                                                                    | NE / —   |
| <i>Trifolium pratense</i> L                             | Fabaceae        | TG1b  | Red clover (forage crop)                  | CH –             | Wild populations of a major forage crop                                                                             | LC / —   |
| <i>Urospermum dalechampii</i> (L.) Scop. ex F.W.Schmidt | Asteraceae      | WHP   | —                                         | GE =             | Preboggion component, cited by 82% of the Genoese sample                                                            | NE / —   |
| <i>Urtica dioica</i> L                                  | Urticaceae      | WHP   | —                                         | GE, CH, CN, MO = | Recorded in all four areas; fibre and leaf-crop candidate                                                           | LC / —   |
| <i>Vaccinium myrtillus</i> L                            | Ericaceae       | TG2–4 | Highbush and lowbush blueberry            | CN =             | Persisting; supported by the berry trade; fruit harvest regulated as an understorey product (Regione Piemonte 1982) | LC / —   |
| <i>Vaccinium vitis-idaea</i> L                          | Ericaceae       | TG1b  | Lingonberry                               | CN –             | Cultivated in northern Europe                                                                                       | LC / —   |
| <i>Valerianella locusta</i> (L.) Laterr                 | Caprifoliaceae  | TG1b  | Corn salad, lamb's lettuce                | CN –             | Wild populations of a European leaf crop                                                                            | LC / —   |

**Table 2** (continued)

| Taxon                                                | Family         | Class | Related crop(s) | Areas; trend | Notes                                         | Red List |
|------------------------------------------------------|----------------|-------|-----------------|--------------|-----------------------------------------------|----------|
| <i>Veronica cf. allionii</i> Vill                    | Plantaginaceae | WHP   | --              | CN –         | High-altitude tea substitute                  | NE / —   |
| <i>Viola odorata</i> L. (and <i>V. calcarata</i> L.) | Violaceae      | WHP   | --              | CN, MO –     | Formerly sold to confectioners and herbalists | LC / —   |

For each taxon, informants were also asked whether the plant was, or had been, transplanted, sown, tended in place, protected from cutting or otherwise deliberately managed. Reports of such management were used to identify cases of incipient domestication, defined as a wild-gathered taxon undergoing a documented transition towards cultivation.

### Statistical analysis

Trends in use were coded from informants' own accounts as increasing or newly commercialised (+), persisting (=), or declining or no longer gathered (–). Each taxon was also assigned to the vegetation belt and habitat in which informants actually gathered it, following the habitat indications given during interviews and field walks and checking them against Flora d'Italia (Pignatti et al. 2017–2019); the assignment is coarse, and several taxa span belts, but it is sufficient for the contrasts examined in Sect. 3.7. The association between vegetation belt and reported trend was tested with Fisher's exact test, using the two-by-two contingency of belt (montane–subalpine and alpine, broadly above about 1500 m, versus hill and lowland) against trend (declining versus not declining). Given the small numbers involved, these tests are reported as descriptive rather than confirmatory.

### Ethics

The research followed the Code of Ethics of the International Society of Ethnobiology (ISE 2006). The Ethics Committee of the University of Gastronomic Sciences, Pollenzo, Italy, granted ethical approval under protocol number UNISG 24092024.

We obtained prior informed consent verbally from every participant before each interview. We explained the research aims, the intended use of the data, and the right to withdraw at any time in the participant's own language in advance, and we requested permission to record separately. No personal data beyond age, occupation, and locality are reported here; informants interviewed in a private capacity are identified by initials or first name only, whereas professional figures who explicitly agreed to be named are cited in full. Free, prior and informed consent covered both the interview itself and the publication of anonymised quotations.

## Results

### The ethnobotanical inventory

Pooling the four datasets and harmonising nomenclature yielded 62 wild taxa across 30 families (Table 1). Asteraceae dominate the inventory with 14 taxa, followed by Rosaceae (6), Lamiaceae (4), and Amaryl-lidaceae and Plantaginaceae (3 each); the remaining families contribute one or two taxa each. This is the expected signature of a European leafy-greens tradition, in which the composite rosettes gathered before flowering carry most of the culinary weight.

The distribution across areas is strikingly uneven. Cuneo contributed the largest repertoire (38 taxa), reflecting both its altitudinal range and a distinct Alpine tradition of aromatic herbal preparations; the Chierese contributed 21, the Genoese area 17, and Monferrato 14. Only two taxa—*Urtica dioica* L. and *Taraxacum officinale*—were recorded in all four areas. Five further taxa were recorded in three areas each: *Borago officinalis* L., *Fragaria vesca* L., *Humulus lupulus*, *Malva sylvestris* L. and *Silene vulgaris* (Moench) Garcke. Forty-three taxa, just under 70% of the inventory, were quoted in a single area only. Given the short distances involved, this is less a matter of flora than of culinary tradition: *Reichardia picroides* and *Urospermum dalechampii* are readily available in Piedmont but are not (anymore?) considered food there. At the same time, the Alpine *Artemisia* complex has no counterpart in the Ligurian repertoire.

Nomenclature and folk taxonomy diverge in instructive ways. In the Genoese sample, 14 botanical species were designated by 17 dialect names: *Scixerboa* and *Cardelle* both denote *Sonchus oleraceus*, *Talegua* and *Gattalevre* both denote *R. picroides*, and *Piscianletto* and *Garatun* both refer to the *Hyoseris radiata* group. Rather than imprecision, this granularity indicates a folk taxonomy fine enough to discriminate among the many edible-leaved Asteraceae that make up the mixture.

### Crop wild relatives and wild harvested plants in the inventory

Applying the taxon group concept to the inventory (Table 2) shows that the foraged repertoire and the national register of plant genetic resources overlap to a degree that is not obvious from either

literature alone. Forty-three of the 62 taxa (69%) are CWR in the broad taxon-group sense. Twenty-eight (45% of the inventory) are conspecific with a cultivated plant and therefore belong to taxon group 1b: they are wild or feral populations of a species that is itself a crop. A further 15 (24%) are congeneric with a crop without being conspecific with it. The remaining 19 taxa (31%) are wild-harvested plants with no cultivated congener.

The taxon group 1b set is not a marginal fraction of the inventory but includes several of its most culturally salient members. *Beta vulgaris* subsp. *maritima* (L.) Arcang., the wild progenitor of all cultivated beets, is one of the six species cited by every informant in the Genoese sample and is a defining component of *Preboggion*. *Humulus lupulus*, the *Luvertin* of the Piedmontese material and the only taxon reported as gathered in undiminished quantity, is the crop species itself. *Corylus avellana* L. is the species on which the Piedmontese hazelnut sector rests, and *Prunus avium* (L.) L. that on which the cherry orchards of the Chierese rest, in both cases with wild and feral populations persisting alongside the orchards. *Asparagus officinalis* L., *Portulaca oleracea* L., *Valerianella locusta* (L.) Laterr., *Rumex acetosa* L., *Nasturtium officinale* W.T.Aiton, *Foeniculum vulgare* Mill., *Sanguisorba minor* Scop., *Ribes rubrum* L., *Juglans regia* L. and *Cornus mas* L. are all wild populations of European crops, several of them of considerable local importance.

The distribution of reported trends across the three classes is informative (Table 3). All five taxa reported as increasing or newly commercialized—*Allium ursinum*, *Asparagus officinalis*, *H. lupulus*, *Matricaria chamomilla* L. and *P. oleracea*—are CWR, and four of the five belong to taxon group 1b. No wild-harvested plant was reported as increasing. Conversely, 12 of 28 taxon group 1b taxa (43%) were reported as declining, compared with 10 of 15 congeneric taxa (67%) and 11 of 19 wild-harvested plants (58%). Neither contrast reaches conventional significance with these numbers (Fisher's exact test,  $p=0.20$  for decline and  $p=0.17$  for increase, taxon group 1b against all others), and we present them as patterns to be explained rather than as demonstrated effects. The direction, however, is consistent: the part of the repertoire that is holding or expanding is composed almost entirely of the wild relatives of cultivated plants, and

**Table 3** Reported trend in use by CWR class. Classes as in Table 2

| Class                         | n  | Increasing (+) | Persisting (=) | Declining (–) |
|-------------------------------|----|----------------|----------------|---------------|
| TG1b: conspecific with a crop | 28 | 4              | 12             | 12            |
| TG2–4: congeneric with a crop | 15 | 1              | 4              | 10            |
| WHP: no cultivated congener   | 19 | 0              | 8              | 11            |
| Total                         | 62 | 5              | 24             | 33            |

**Fig. 4** *Gentiana lutea* L. in flower in a montane meadow of the province of Cuneo; the roots are the part gathered (Photo: A. Pieroni)

the part that is disappearing fastest is the part with no cultivated counterpart.

Three cases in the inventory document an ongoing transition from gathering towards cultivation. In the Racconigi plain, *M. chamomilla* has shifted within living memory from a field weed collected opportunistically to a plant deliberately transplanted and grown. *Gentiana lutea* L. (Fig. 4) and *Artemisia umbelliformis* Lam., the two Alpine taxa whose harvest supports the gentian and *Genepi* herbal-preparation trades, have both moved partly into cultivation drawn from wild germplasm. Alongside

these, informants in three areas reported transplanting *F. vesca* and *R. rubrum* seedlings from woodland and hedgerow into kitchen gardens, a management practice short of cultivation but well beyond simple harvesting.

None of the 62 taxa is of conservation concern at the species level. Where a Red List assessment exists, at the global or the European level, the category is invariably Least Concern. The only recorded taxon listed above that level in the national Red List of the Italian flora is *Gentiana lutea*, classified as Near Threatened in Italy owing to the commercial collection of its roots (Bilz et al. 2011; Allen et al. 2014; Orsenigo et al. 2021; IUCN 2026); the category of each taxon at both levels is reported in Table 2. None of the recorded taxa figures in the list of absolutely protected species annexed to the Piedmontese law, although *Gentiana lutea* and *Vaccinium myrtillus* L. enjoy absolute protection in single provinces (Alessandria and Asti, respectively); the gathering of the three *Genepi* species and of *G. lutea* roots falls under the authorisation regime for wild officinal plants, and *Vaccinium* fruits are harvested as regulated understorey products (Regione Piemonte 1982). This is consistent with the profile of the inventory as a whole: what is gathered is, almost without exception, the common flora of frequented habitats, and the conservation questions raised in the Discussion concern populations, practices and habitats rather than species survival.

**Preboggion:** a mixture that outlives its species

In the Genoese sample, all eleven informants knew and practised *Preboggion* gathering, with an average of some sixty years of experience. Every informant cited six species—*B. officinalis*, *B. vulgaris* subsp. *maritima*, *T. officinale*, *S. vulgaris*, *S. minor* and *U. dioica*—which together constitute the identity core of the mixture; *M. sylvestris* (91%) and *U. dalechampii*

(82%) follow closely. Informants consistently described the mixture as variable by season, slope and gatherer, and treated that variability as constitutive rather than accidental.

The comparison of past and present practice reported by the same informants is unambiguous. Frequent use of *Preboggion* fell from 93 to 34%; use in soups fell from 87 to 28%; use in savoury pies held up better, falling from 80 to 41%; and family transmission of the knowledge fell from 100 to 38%. Averaged across indicators, the decline is roughly by half. The most consequential figure, however, is the last: the species themselves are common, resilient, and in no sense threatened, but the capacity to read them as food is not being handed on. As one informant put it, people are now afraid of getting the identification wrong.

What keeps *Preboggion* alive is precisely that it is a named artefact with a market. At the Mercato Orientale it is sold ready-mixed and correctly composed; the buyer purchases not only the greens, but the identification skill embedded in them. This is the clearest instance in our material of a wider pattern: where gathering knowledge has been externalised into a commodity, consumption survives the collapse of household transmission.

#### Luvertin: the resilient shoot

*Humulus lupulus* is the only taxon in our material that all Piedmontese informants described as still being gathered in essentially unchanged quantities, and the reasons given are consistent across the three areas: the shoots are easy to recognise, they appear in a narrow and well-known season, and they are unambiguously good to eat. The plant carries at least three folk names in the Monferrato alone (*Luvertin*, *Uvertin*, *Invertin*), it supports a local fair at Sant'Albano Stura, and it has moved into professional kitchens—as a filling for fresh pasta in the everyday restaurant sector, and as a seasonal signature in the fine-dining sector, where the chef interviewed in the Monferrato described sourcing and handling as the decisive steps.

The same species also illustrates how a single plant can span registers. Its female cones are used domestically as a bitter tonic and sleep aid, in doses reported to us in the same terms as those given in mid-twentieth-century herbal manuals, and industrially in brewing. Very few of the taxa in Table 1 travel this

comfortably between the kitchen garden, the pharmacy shelf and the restaurant. It is worth noting that the wild populations gathered for *Luvertin* and the plants grown in the region's small but expanding hop yards are the same species, so that the shoot harvest and the cone crop draw on a single gene pool.

#### Erosion of the transmission chain

In all four areas, informants located the break in transmission in the same period: the post-war decades and, above all, the economic boom of the 1960s, with rural depopulation, the arrival of industrial foodstuffs perceived as more prestigious, and the association of wild greens with poverty. The Occitan formulation recorded in Cuneo is the sharpest statement of this reversal: “Nous sen pas ciabre”—we are not goats. Eating what the mountain offered became embarrassing just as it stopped being necessary.

Older informants described learning in the field, from mothers and grandmothers, as small children carrying a basket. That channel is now largely closed. In the Chierese, a nursery grower observed that the habit has gone and the memory with it, while his wife added that people no longer know how to look because the word is no longer passed from one generation to the next. A young farmer in the same area asked whether he could recognise Coujet (*S. vulgaris*) in a meadow and admitted he would struggle. The Genoese figures quantify the same phenomenon: perceived knowledge of wild greens among younger generations stands at 45% against 100% for the previous ones.

Alongside the social causes, informants named ecological ones: agricultural intensification and specialisation, heavy machinery and soil compaction, herbicides, the loss of field hedges, the disappearance of permanent meadows to building and to arable conversion, and, at altitude, grazing pressure on *Vaccinium* stands and the effect of harvesting combs that strip leaves along with berries. Climate was raised repeatedly and specifically: in the Varaita valley, *Genepi* is now in flower by mid-July and already over by August, when gathering traditionally took place.

#### Re-emergence and new actors

The same interviews record a countermovement. Wild hop shoots, wild asparagus, purslane, dandelion

and poppy rosettes now appear on farmers' market stalls in the Chierese, bought by consumers who do not gather. Agriturismo and mountain refuges in Cuneo have built menus around wild plants; several were founded by returnees who left for the city and came back. Piedmont has organised courses on identification and processing, and university-linked projects have re-engaged younger participants with the older documentary record. Informants repeatedly proposed ecomuseums and slow-tourism itineraries as a way forward, and in Liguria the community around the Mercato Orientale is actively pursuing *Preboggion*'s candidacy as a Slow Food Presidium. A Presidium is a scheme run by the Slow Food Foundation for Biodiversity that supports a food product judged to be at risk of disappearing by constituting its producers as a group, agreeing a production protocol, providing technical assistance, and opening commercial outlets.

These initiatives were not described uncritically. Informants in the Cuneo valleys reported persistent consumer distrust of wild ingredients in restaurants and fear of poisoning; the agronomist interviewed in the Chierese warned explicitly against the local proverb that any herb that raises its head makes good soup, insisting that renewed interest is welcome only if matched by competence.

#### Elevational patterning of reported trends

The question is simple: does the reported decline in gathering strengthen at high altitude than in the lowlands? To answer it, we divided the 62 taxa into the two belt groups defined in Sect. 2.5: a montane–subalpine and alpine group of 12 taxa, all of them recorded in the province of Cuneo, and a hill and lowland group of 50 taxa. The two groups behave differently. In the high-altitude group, eight of the 12 taxa (67%) were reported as declining or no longer gathered, four as persisting, and none as increasing. In the hill and lowland group, decline affects exactly half of the taxa (25 of 50, 50%), 20 taxa were reported as persisting, and five (10%) as increasing or newly commercialised; every taxon with a positive trend in the whole inventory therefore belongs to the lower belts. Because Cuneo is the only study area that spans both belts, it also provides an internal control that removes differences between areas: within Cuneo alone the contrast points the same way, with eight of 12 high-belt taxa declining against 11 of 26

lower-belt taxa, three of which are increasing. With samples of this size the association does not reach statistical significance (Fisher's exact test,  $p=0.35$  for the full inventory and  $p=0.30$  within Cuneo), so we read these figures as a consistent descriptive pattern in need of explanation rather than as a demonstrated effect; its possible causes are examined in Sect. 4.4.

## Discussion

### Contraction of breadth, expansion of meaning

Read together, the four datasets do not support either of the two standard narratives. Traditional foraging in NW Italy is not simply disappearing: several taxa are gathered as much as ever, two preparations have acquired a market, and interest among younger and urban consumers is real. Nor is it thriving: the number of taxa any given person can name and use has fallen sharply, entire use categories—famine foods, coffee surrogates, veterinary and household use—have gone, and the channel through which the knowledge moved has been cut. What we observe is a decoupling. The repertoire contracts in breadth while the cultural and commercial value of what remains expands. The same decoupling has been described for the Southern Occitan Alps and the upper Susa Valley (Alrhoun et al. 2025b; Sulaiman et al. 2024b) and, in more advanced form, in coastal Mediterranean settings where the tradition survives mainly as memory (Sulaiman et al. 2026).

Our material suggests what determines which taxa survive the contraction. Three anchors recur. The first is incorporation into a named dish or product: *Preboggion* in Liguria, *Luvertin* in Piedmont, *Genepi* and *Genziana* herbal preparations in the Alpine valleys. A plant that is an ingredient of something with a name is remembered, sought and bought; a plant that is merely edible is forgotten. The second is ease and safety of recognition, which informants stated explicitly for hop shoots and elder flowers. The third is the existence of a commercial channel, whether a market stall, a fair, a herbal-processing workshop or a restaurant. Taxa lacking all three—*Papaver rhoeas* L., *Tragopogon pratensis* L., *Rumex alpinus* L., *Blitum bonus-henricus* (L.) Rchb.—are precisely those our informants named as no longer gathered, and they are

disproportionately the plants that were most closely bound to scarcity.

#### What foraging knowledge contributes to work on CWR

The central result of this study is the extent of the overlap reported in Sect. 3.2: 70% of a repertoire assembled purely on ethnobotanical criteria consists of CWR in the taxon-group sense, and 46% of wild populations of crop species themselves. The overlap is not a coincidence of classification. People gather, preferentially, the wild counterparts of the plants they already grow and eat, because those are the plants whose edibility, seasonality and processing they already understand. Wild food plant inventories and CWR checklists are, to a substantial degree, two descriptions of the same set of populations arrived at from opposite directions: one from use, the other from taxonomy. That largely separate research communities compile them is a historical accident with practical costs.

Those costs are worth stating concretely, because the ethnobotanical record supplies three things that a taxonomic and spatial inventory cannot.

The first is the location and condition of populations that are actually being used. A CWR checklist establishes that *B. vulgaris* subsp. *maritima* occurs on the Ligurian coast; Landucci et al. (2014) place the genus *Beta* among the priority Italian CWR, and it is among the highest-priority genera globally (Vincent et al. 2013; Castañeda-Alvarez et al. 2016). Our informants add that specific stands on the terraced slopes above Bavari are harvested repeatedly throughout the season by named gatherers, that the harvest supplies a specific urban market, and that the people concerned can walk to the stands. That matters, because sea beet is not an abstract genetic resource: it is the population from which the rhizomania resistance locus *Rz2* was mapped directly, by sequencing wild beets sampled in their natural habitat. The authors of that work explicitly concluded that in situ preservation of such populations makes the approach possible (Capistrano-Gossman et al. 2017). A community that harvests a sea beet population weekly is, whether or not it is described in these terms, maintaining an in situ genetic resource and monitoring it continuously. When the harvesting stops, the

monitoring stops with it, and the population becomes a dot on a distribution map again.

The second is phenotypic resolution. Folk taxonomies are frequently finer than the botanical species where the distinctions matter to the user, and coarser where they do not. Our Genoese material shows both: 14 species carry 17 names, while elsewhere a single name covers two species. The names that multiply are those of the edible-leaved Asteraceae that must be discriminated in the field at rosette stage, and the criteria informants gave for the distinctions—bitterness, leaf texture, and the moment at which the rosette becomes unusable—are recognisably the same descriptors that appear in germplasm characterisation. Something similar has been argued for the Basque Country, where this journal documented the fine-grained naming of gathered wild vegetables (Menendez-Baceta et al. 2012). A folk taxonomy of this kind is the sedimented outcome of many generations of phenotypic screening across a wild population, and it is the sort of information that is expensive to reconstruct once lost and cheap to record while it survives.

The third is a defensible shortlist for neo-domestication. The standard triage for identifying wild food plants that might become novel crops combines cultural salience, existing economic value in local markets, and agronomic feasibility (Sulaiman et al. 2023). Ethnobotanical fieldwork supplies the first two directly and cheaply. In our inventory, *R. picroides*, *S. vulgaris*, *S. minor*, *U. dalechampii*, *A. ursinum*, and *P. oleracea* satisfy both criteria: high salience, an existing point of sale or restaurant demand, and several have already attracted attention as candidate leafy vegetables elsewhere in southern Europe. *Humulus lupulus* unusually satisfies them, since the crop already exists and only the wild shoot harvest, rather than the plant, would need to be brought into production.

Beyond candidate identification, our material includes four documented cases of a transition already underway (Sect. 3.2). Domestication is best understood as a process rather than an event, and the early stages—deliberate transplanting, protection in place, movement from field margin to seedbed—are ordinary and unspectacular (von Wettberg et al. 2020). The Racconigi chamomile and the Pancalieri mint are exactly this: plants that entered the regional agricultural economy from the wild flora

within living memory, through the intermediate step of being tended where they stood. Transplanting *F. vesca* and *R. rubrum* into kitchen gardens is the same process caught earlier. None of these transitions is recoverable from a herbarium or a distribution model. They are visible only because someone asked the people doing them, and in three of the four cases the people who could describe the transition were over seventy-five.

The erosion has a genetic resources cost, not only a cultural one

If the argument above holds, then the decline documented in Sects. 3.3 and 3.5 is not only a loss of intangible heritage. Three distinct mechanisms connect it to the state of the genetic resources themselves.

The first operates through habitat. Several declining taxa are not merely found in a habitat but depend on a management regime that included gathering. *Rumex alpinus* and *B. bonus-henricus* are nitrophilous species of livestock resting places, of the surroundings of alpine cowsheds and of seasonally inhabited high settlements; their habitat is a by-product of transhumant husbandry, and the same is largely true of the moist mown meadows in which *Allium schoenoprasum* L. was gathered. When the summer pastoral economy contracts, the stands contract with it, so that the plant and the gatherer disappear in the same movement and for the same reason. This mechanism is repeatedly documented for Alpine grassland flora following the withdrawal of traditional management (Tasser et al. 2007; Niedrist et al. 2009). Here the loss of practice and the loss of population are not two events but one.

The second operates through land use, and affects a well-defined guild: the plants of hay meadows and of low-input arable fields and their margins. *P. rhoeas*, *Sinapis arvensis* L., *V. locusta*, *T. pratensis*, *Salvia pratensis* L., *Trifolium pratense* L., *Bellis perennis* L., *R. acetosa*, *Plantago lanceolata* L., *Leontodon hispidus* L. and, at the montane meadow margin, *Primula veris* L. are precisely the guilds whose measured decline across Europe has been attributed to fertiliser and herbicide use and to the conversion of permanent grassland (Storkey et al. 2012), and in the Alpine context to the parallel intensification of accessible land and abandonment of the rest (Tasser et al. 2007; Niedrist et al. 2009). Our informants named the

same drivers independently and without prompting. It is worth noting how many members of this guild are CWR: *V. locusta* is the crop species itself, *T. pratense* is a major forage crop, *R. acetosa* is cultivated sorrel, and *T. pratensis* and *S. arvensis* are congeners of salsify and of the *Brassica* crops respectively. The arable and meadow margin is not an ethnobotanical curiosity but one of the principal reservoirs of European CWR diversity, and it is being lost for reasons unrelated to whether anyone still gathers from it.

The third operates through knowledge, and is the one specific to this study. Where a population persists and its habitat persists, what can still be lost is the ability to find it, to recognise it, and to distinguish what is worth taking within it. This is the loss our Genoese figures measure most directly: the species are common and unthreatened, family transmission has fallen from 100 to 38%, and informants cited fear of misidentification. Nothing about the beet populations above Genoa has changed; what has changed is how many people can read them. Recent ethnobiological work has argued that local ecological knowledge cannot be maintained in the absence of continuous, embodied interaction with the landscape (Pieroni et al. 2025; Sulaiman et al. 2026), and our material is consistent with that: the taxa that hold or expand are, almost without exception, those of habitats crossed daily anyway: hedgerows and riparian scrub for *H. lupulus*, vineyard and field margins for *A. officinalis*, gardens and yards for *P. oleracea*, *Melissa officinalis* L. and *M. chamomilla*, roadsides and orchard floors for *T. officinale*, and the terraced slopes immediately above Genoa for the whole *Preboggion* assemblage. Gathering survives where it costs no additional journey. What has contracted at altitude is not the flora but routine bodily presence in the habitat, the food- and landscape-specific form of what has been described more generally as the extinction of experience (Soga and Gaston 2016).

Elevation, frequentation and an attribution problem

The elevational patterning reported in Sect. 3.7 deserves a closer look, because it bears on where in situ measures would be effective.

The obvious reading is that people no longer go up. Against this, however, stands an internal control that we consider decisive. In the same belt, on the same slopes, and in several cases in the same valleys, four

taxa are reported as fully persisting: *A. umbelliformis*, *Achillea erba-rotta* All. subsp. *moschata* (Wulfen) I. Richardson, *G. lutea* and *Vaccinium myrtillus*. Each of these four has a current commercial outlet—the *Genepi* and gentian herbal preparations, and the berry trade. Each of the nine that are disappearing lacks one, and the one that once had one, *Pinus cembra*, lost it when that trade closed. Whatever is removing the nine cannot be a property of the high belt as such, because it leaves the four untouched. Within the alpine belt, availability is not the discriminating variable; the existence of a reason to climb is.

What, then, of climate? Our informants raised it specifically, but more narrowly than the question is usually posed. In the Varaita valley, they reported that *Genepi* is now in flower by mid-July and already over by August, and at altitude they described grazing pressure on *Vaccinium* stands, together with damage from harvesting combs. Both observations are consistent with the wider evidence. Vascular plants in the Alps have shifted upward measurably over recent decades (Lenoir et al. 2008; Parolo and Rossi 2008), summit floras have been enriched at an accelerating rate that tracks warming (Pauli et al. 2012; Steinbauer et al. 2018), and range dynamics are slower at the highest elevations, so that alpine specialists lag behind the climate they experience (Rumpf et al. 2018). Yet three things argue against reading our declines as a climate signal. None of the nine retreating taxa is rare or threatened in Piedmont; the documented direction of change on Alpine summits is enrichment rather than the loss of common alpine species; and, again, climate cannot explain why four taxa of the same belt are stable. The defensible reading is narrower and, we think, more interesting: climate is acting on our system as a modifier of the gathering calendar rather than as a cause of the loss of the practice. A phenological advance of two to three weeks is a minor inconvenience for someone living on the mountain and a disqualifying obstacle for someone who now drives up on a fixed summer weekend.

This points to an attribution problem that deserves to be stated plainly, because it affects the whole diachronic ethnobotanical literature and, by extension, any use of that literature to infer the status of wild populations. Abandonment and warming produce closely similar landscape signals: the canonical demonstration is that most of the recent advance

of the tree line in the Swiss Alps represents infilling of abandoned grassland rather than a true climatically driven upward shift (Gehrig-Fasel et al. 2007). Informants who report that there is less of a plant may be reporting a reduced encounter rate rather than a reduced abundance, and reduced encounter follows from reduced presence as surely as from reduced population. Our data, like almost all data of this kind, measure use and encounter, not abundance. Testing the alternatives properly would require paired vegetation surveys in the abandoned and still-grazed sectors of the same valleys, which we did not carry out. For CWR assessment, this is an important caveat: ethnobotanical data are excellent evidence of use, management, and location, but weak evidence of population size.

#### Commercialisation as a double-edged conservation instrument

Section 4.4 established that a commercial outlet is what keeps a high-belt taxon in use. Section 3.2 established that every taxon in the inventory with a positive trend is a CWR. Together these results suggest that the market is currently the most effective instrument maintaining wild plant use in the region—and that it is a hazardous one.

Three implications follow. First, commercialisation can conserve practice while hollowing out competence, and the two outcomes should be evaluated separately. The Mercato Orientale keeps *Preboggion* in circulation, and in doing so it allows the buyer to acquire the mixture without acquiring the knowledge that composed it. Persistence of use and persistence of transmission are different outcomes, and our data show the first improving while the second continues to decline. This is consistent with a finding from a large Iberian dataset: the wild plant uses that persist are anchored to high cultural appreciation and the recreational value of gathering rather than necessity (Reyes-García et al. 2015).

Second, demand concentrated on a few charismatic taxa creates harvesting pressure that traditional dispersed gathering never generated, and the ecological effects of concentrated wild harvest are well documented and frequently underestimated (Ticktin 2004). The Cuneo material already contains the historical precedent in its own record: the early twentieth-century bulk trade in *Artemisia* for the aromatic-herb

industry stripped several species from the valleys. That episode is, in the vocabulary of this journal, a documented case of genetic erosion of a wild harvested resource driven by commercial demand, and it is remembered locally. It is worth stating explicitly, because the point bears directly on the conservation implications of the practice, that no informant in any of the four areas attributed the decline of a currently gathered taxon to gathering pressure itself: the drivers named were land-use change, the contraction of the pastoral economy and the failure of transmission (Sects. 3.5 and 3.7). The two partial exceptions in our material are historical or marginal—the *Artemisia* episode just described, and the damage done to *Vaccinium* stands by harvesting combs, which informants linked to the commercial berry trade rather than to domestic gathering. The risk is therefore prospective: it would arise precisely where the revival documented in Sect. 3.6 concentrated demand on wild stands without the parallel movement into cultivation observed for *Gentiana lutea* and *Artemisia umbelliformis*. Access and harvest rules strongly condition what can actually be gathered in European settings (Mina et al. 2023). Current Piedmontese law (Regional Law no. 32 of 2 November 1982; Regione Piemonte 1982) prohibits outright the collection of an annexed list of absolutely protected species, allows for the remaining wild flora a daily harvest of five specimens per person without uprooting of the underground organs, exempts the most commonly consumed edible species from that limit, and provides for individual derogations for residents for whom collection is a source of seasonal income; in Liguria, wild flora protection rests on Regional Law no. 9 of 30 January 1984, whose species annexes were replaced and updated by the regional biodiversity law of 2009, which distinguishes fully protected species from species whose gathering is regulated (Regione Liguria 2009). An apparatus of this kind is difficult to reconcile with commercial supply and was not designed for a revival of the sort described here. Any policy that deliberately creates a reason to climb—and Sect. 4.4 implies that this is the only measure likely to recover high-belt use—must be paired with harvest regulation designed for the purpose. Nor is the tension peculiar to Italy. In Israel and Palestine, long-standing protection orders on za'atar (*Origanum syriacum* L.) and 'akkoub (*Gundelia tournefortii* L.) have been credited with relieving pressure on wild

populations while simultaneously criminalising a traditional, largely Palestinian, foraging practice, and they have become an instructive case of how collection restrictions redistribute both ecological pressure and cultural loss (Harris 2023; Braverman 2024). The general lesson applies here: harvest regulation is never only a conservation instrument, since it also determines who may continue to exercise, and therefore to transmit, the knowledge that Sect. 4.3 identified as the resource actually at risk.

Third, formal recognition schemes face a genuine dilemma that has a direct genetic resources dimension. A Slow Food Presidium would give *Preboggion* visibility, protect its local cultivars and support its gatherers. Food festivals and similar arenas have been shown to function effectively as settings in which dormant food knowledge is re-performed and re-legitimated (Fontefrancesco and Zocchi 2020). Still, every recognition scheme tends to fix a composition, and the variability of *Preboggion*—by season, slope and gatherer—is not a defect to be corrected but the very form the knowledge takes, and arguably a proxy for the diversity of the populations being harvested. A protocol that fixed a canonical species list would standardise the harvest onto a subset of the stands. The heritagisation literature warns, more generally, that such arenas select which elements of a foodscape become recognised heritage and which actors are authorised to speak for it (Zocchi et al. 2021). The most useful design would specify a protected core, a permitted set, and an explicit right to seasonal and local variation.

#### Who reinvents foraging, and why it matters for conservation

A design feature of this study turns out to be one of its more informative results. We did not sample the population that the international literature treats as the protagonist of the contemporary foraging revival. Urban and peri-urban recreational foragers have been described in detail in Berlin (Landor-Yamagata et al. 2018), Vienna (Schunko et al. 2021), Baltimore (Synk et al. 2017), and North American cities (Poe et al. 2013; McLain et al. 2014), and the practice is documented across a wide range of urban contexts (Shackleton et al. 2017; Fischer and Kowarik 2020); their motivations, largely recreational, health-related and provenance-driven,

are well characterised (Schunko et al. 2015; Reyes-García et al. 2015). None of these people is in our sample. Our informants were rural and peri-urban residents, most over sixty-five, selected for their long-term and inherited experience of gathering. And yet innovation is unmistakably present in what they described.

Five figures recur across the four areas: the returnee producer, who converts a domestic seasonal practice into a bookable service and in doing so needs the plants to be reliably available and nameable (Dematteis 2011; Corrado et al. 2014); the market vendor, who standardises the mixture just enough for it to be sold; the chef, for whom wild plants are a seasonal signature and whose knowledge comes from books, suppliers and peers rather than from a family line, exactly as documented for nearby Lombardy (Sulaiman et al. 2024a); the institutional broker behind a Presidium candidacy, a fair or an ecomuseum; and the custodian turned teacher, who learned in the field as a child and now transmits to strangers rather than to kin. None is a hobbyist: each has an economic or institutional stake in the plants, and all five are intermediaries standing between the plant and a public that does not gather.

Two features of this configuration matter for genetic resources. The innovation is organisational rather than botanical: nothing in Table 1 is new, no taxon entered the repertoire through the contemporary revival, and the species with a positive trend are long-standing local foods moving through new formats. This contrasts usefully with urban foraging inventories, which are typically floristically broader and include ornamental, alien, and spontaneously urban taxa absent from any inherited local corpus (Landor-Yamagata et al. 2018; Fischer and Kowarik 2020). And the innovators are, in general, not the transmitters: a vendor who sells a correctly composed mixture, a chef who buys shoots from a supplier and a festival that stages a plant once a year all keep the plant in circulation without restoring in-field learning. For anyone hoping to use these actors as custodians of in situ genetic resources—and they are the obvious candidates, since they have a stake in the populations continuing to exist—this distinction is crucial. They can be relied upon to maintain demand. They cannot, as currently organised, be relied upon to maintain the competence that locates and discriminates the populations.

A further observation, which our sampling allows us to state but not to quantify, concerns gender. The transmission that broke was largely female and domestic, and the knowledge holders we interviewed in Cuneo were predominantly women. The reinvention is being carried out to a considerable extent through commercial and public roles—restaurateurs, herbal processors, market vendors, fair committees—in which men are, at least in our sample, more visible. If confirmed, this would mean that the shift from household to market has also shifted who is credited with the knowledge, a dynamic the food heritage literature identifies as a recurrent risk of heritagisation (Zocchi et al. 2021).

### Recommendations

Three proposals follow from the material, addressed to different audiences.

To those compiling national inventories of plant genetic resources, we suggest treating ethnobotanical surveys as a routine input rather than as parallel literature. Concretely, the Italian CWR and wild-harvested plant checklist (Landucci et al. 2014; Perrino and Perrino 2020) could carry a use-and-management attribute for each taxon, recording whether the taxon is currently gathered, whether it is deliberately managed, and whether a community exists that can locate and discriminate its populations. That attribute is cheap to collect alongside existing ethnobotanical fieldwork, it changes the priority ranking, and it identifies the populations for which in situ measures have a realistic chance of being maintained by the people already present.

To those designing in situ measures in this region specifically, the asymmetry established in Sect. 4.4 has a clear consequence. Where loss is driven by access rather than availability—the alpine taxa—measures that restore a reason to be present, such as grazing support, staffed refuges, guided harvest, and short product chains, will recover use faster than measures that protect plants that are in no danger. Where loss is driven by habitat conversion—the meadow and arable margin guild, which contains a disproportionate share of the region's CWR—the priority is the reverse, and no amount of cultural interest will bring *V. locusta* or *P. rhoëas* back onto a plate without margins, hedgerows and low-input permanent grassland. The terraced slopes above Genoa are a

candidate for the first kind of measure and, given the identity of the species involved, a plausible site for a genetic reserve built around a harvesting community rather than around an exclusion.

For those documenting the knowledge, the practical point is sequence. Systematic documentation is most urgent where the last competent gatherers are now in their eighties, which in our material means the Occitan valleys and the Genoese hinterland. Recognition schemes should protect variability rather than fix a recipe. And the single most consequential intervention, on our evidence, is also the least technical: arrangements that put learners and knowledge holders in the same field in the same season. No amount of documentation reproduces a practice that has stopped being practised, and no distribution map substitutes for a person who knows which slope.

### Limitations

The four datasets were collected independently, with partly different interview schedules and sampling strategies, and the samples are small and purposive; the frequencies reported for the Genoese area in particular rest on eleven informants and should be read as indicative rather than representative. Fieldwork was concentrated in late winter and spring, which favours leafy species and under-records autumn taxa. Perceived past frequencies of use are retrospective self-reports and are subject to the usual recall effects, including a tendency to contrast a vivid childhood with an undifferentiated present. Our samples are strongly skewed towards older residents, and the perspective of younger non-gatherers is represented only through the accounts of others; for the same reason, the study contains no urban recreational foragers, and the contemporary urban demand described in Sects. 4.5 and 4.6 is documented only from the supply side.

Two limitations bear specifically on the genetic resources argument. We collected no vegetation or abundance data, so the ecological reading proposed in Sect. 4.4 rests on reported use and the published literature rather than measured changes in plant populations; as noted, ethnobotanical data provide strong evidence of use and location and weak evidence of population size. And the CWR assignment in Table 2 rests on the taxon group concept rather than on documented crossing behaviour or molecular data

(Maxted et al. 2006). For the taxon group 1b taxa, this is unproblematic, since conspecificity with the crop is not in doubt; for several congeneric assignments, the relationship with the crop gene pool is remote, and the corresponding figures should be read as an upper bound on the breeding-relevant fraction of the inventory.

### Conclusion

Across four areas of Liguria and Piedmont, 62 wild taxa in 30 families were recorded as gathered now or within living memory, and only two of them are shared by all four areas. Just under seventy per cent of that repertoire consists of CWR in the taxon group sense, and 45% of wild populations of crop species themselves, including the wild progenitor of all cultivated beets, which is the identity core of a mixture still sold daily in the centre of Genoa. Every taxon in the inventory whose use is expanding is a CWR; not one wild harvested plant without a cultivated counterpart is gaining ground.

Behind the low overlap between areas lies a general pattern: what survives is not the plant but the artefact the mixture, the shoot, the aromatic preparation and the market or the table that keeps it in circulation. Gathering has ceased to be a household competence transmitted in the field and is becoming a specialised, purchasable and increasingly professional one. Almost every taxon in Table 1 is common, and species conservation is not the issue. What is at issue is maintaining the practices that keep specific populations under continuous local management and observation, and the knowledge that locates and distinguishes them.

We therefore suggest treating the ethnobotany of wild food plants not as a parallel literature to the conservation of plant genetic resources but as one of its instruments. It identifies which wild relatives are embedded in living agroecosystems rather than merely present in them; it supplies phenotypic and locational detail that taxonomic inventories cannot; it catches incipient domestication in progress; and it names the communities on whose continued presence in situ measures would depend. It also has a short window. In our material, people over seventy-five described three of the four documented transitions from gathering towards cultivation.

**Acknowledgements** Special thanks go to all study participants for agreeing to share their local plant knowledge and wisdom, welcoming the researchers into their homes, kitchens, gardens, and meadows, and sharing knowledge that, in many cases, they had not been asked about for decades. We warmly thank Maria Clotilde Giordano (Chierese), Alessia Oderda and Aurora Romano (Province of Cuneo), Ludovica Imarisio (Monferrato), and Francesco Assembri, Lorenzo Fierro, Giovanni Frontani and Francesco Lamorica (Genoa and Bavaria) for having conducted some parts of the fieldwork. We also thank Dr Dauro Mattia Zocchi, University of Bergamo, Italy, who passed away in spring 2026, for his inspiring insight into the initial idea of this work.

**Author contributions** Conceptualisation: AP; Methodology: AP; Investigation: AP; Formal analysis: AP, AKM; Writing—original draft preparation: AP; Writing—review and editing: AP, AKM. All authors read and approved the final manuscript.

**Funding** The authors declare that no funds, grants or other support were received during the preparation of this manuscript.

**Data availability** The inventory supporting the conclusions of this article is included within the article as Tables 1 and 2. The interview transcripts and field notes are not publicly available because they contain information that could compromise participants' privacy.

## Declarations

**Conflict of interests** The authors have no relevant financial or non-financial interests to disclose.

**Ethics approval** The research was conducted in accordance with the Code of Ethics of the International Society of Ethnobiology (ISE 2006). The Ethics Committee of the University of Gastronomic Sciences, Pollenzo, Italy granted ethical approval (protocol UNISG 24092024).

**Consent to participate** We obtained prior free and informed consent verbally from each participant before each interview, after explaining the aims of the research, the intended use of the data, and the possibility of withdrawing at any time in the participant's own language. Permission to record was requested separately.

**Consent for publication** All participants consented to the publication of the anonymised data and of the quotations reproduced here. We report no personal data beyond age, occupation, and locality: informants interviewed in a private capacity are identified by initials or first name only, whereas professional figures who explicitly agreed to be named are cited in full.

**Plant material** No plant material was collected, exported, or otherwise utilised during this study; therefore, no access and benefit sharing procedure under the Nagoya Protocol was applicable. Voucher specimens consulted for verification were collected during previous expeditions by the same research group and are deposited as stated in Sect. 2.3.

## References

- Allen D, Bilz M, Leaman DJ, Miller RM, Timoshyna A, Window J (2014) European red list of medicinal plants. Publications Office of the European Union, Luxembourg
- Alrhoun M, Guiggi V, Gillani SW, Manzoor M, Sulaiman N, Pieroni A (2025a) Diachronic changes in local food heritage: the ethnobiology of wild foods in central Tuscany. *J Ethnobiol Ethnomed* 21:48. <https://doi.org/10.1186/s13002-025-00797-x>
- Alrhoun M, Sulaiman N, Pieroni A (2025b) Shifting herbal knowledge: the ecological and cultural dynamics behind plant use changes in the Southern Occitan Alps. *Plants* 14:367. <https://doi.org/10.3390/plants14030367>
- Alrhoun M, Romano A, Sulaiman N, Pieroni A (2025c) Old plants for new food products? The diachronic human ecology of wild herbs in the western Alps. *Plants* 14:122. <https://doi.org/10.3390/plants14010122>
- Bharucha Z, Pretty J (2010) The roles and values of wild foods in agricultural systems. *Philos Trans R Soc B* 365:2913–2926. <https://doi.org/10.1098/rstb.2010.0123>
- Bilz M, Kell SP, Maxted N, Lansdown RV (2011) European red list of vascular plants. Publications Office of the European Union, Luxembourg
- Braverman I (2024) Green gold: the Akkoub's settler ecologies. *LA+ Interdiscip J Landsc Archit* 19:38–45
- Capistrano-Gossmann GG, Ries D, Holtgräwe D, Minoche A, Kraft T, Frerichmann SLM, Rosleff Soerensen T, Dohm JC, González I, Schilhabel M, Varrelmann M, Tschöep H, Uphoff H, Schütze K, Borchardt D, Toerjek O, Mechelke W, Lein JC, Schechert AW, Frese L, Himmelbauer H, Weisshaar B, Kopisch-Obuch FJ (2017) Crop wild relative populations of *Beta vulgaris* allow direct mapping of agronomically important genes. *Nat Commun* 8:15708. <https://doi.org/10.1038/ncomms15708>
- Castañeda-Álvarez NP, Khoury CK, Achicanoy HA, Bernau V, Dempewolf H, Eastwood RJ, Guarino L, Harker RH, Jarvis A, Maxted N, Müller JV, Ramirez-Villegas J, Sosa CC, Struik PC, Vincent H, Toll J (2016) Global conservation priorities for crop wild relatives. *Nat Plants* 2:16022. <https://doi.org/10.1038/nplants.2016.22>
- Cornara L, La Rocca A, Marsili S, Mariotti MG (2009) Traditional uses of plants in the Eastern Riviera (Liguria, Italy). *J Ethnopharmacol* 125:16–30. <https://doi.org/10.1016/j.jep.2009.06.021>
- Dempewolf H, Baute G, Anderson J, Kilian B, Smith C, Guarino L (2017) Past and future use of wild relatives in crop breeding. *Crop Sci* 57:1070–1082. <https://doi.org/10.2135/cropsci2016.10.0885>
- Corrado F, Dematteis G, Di Gioia A (eds) (2014) Nuovi montanari. Abitare le Alpi nel XXI secolo. FrancoAngeli, Milan
- Fischer LK, Kowarik I (2020) Connecting people to biodiversity in cities of tomorrow: is urban foraging a powerful tool? *Ecol Indic* 112:106087. <https://doi.org/10.1016/j.ecolind.2020.106087>
- Fontefrancesco MF, Zocchi DM (2020) Reviving traditional food knowledge through food festivals. The case of the pink asparagus festival in Mezzago Italy. *Front Sustain*

- Food Syst 4:596028. <https://doi.org/10.3389/fsufs.2020.596028>
- Fontefrancesco MF, Zocchi DM, Cevasco R, Dossche R, Abidullah S, Pieroni A (2022) Crumbotti and rose petals in a ghost mountain valley: foraging, landscape, and their transformations in the upper Borbera Valley, NW Italy. *J Ethnobiol Ethnomed* 18:42. <https://doi.org/10.1186/s13002-022-00535-7>
- Dematteis G (ed) (2011) Montanari per scelta. Indizi di rinascita nella montagna piemontese. FrancoAngeli, Milan
- Gehrig-Fasel J, Guisan A, Zimmermann NE (2007) Tree line shifts in the Swiss Alps: climate change or land abandonment? *J Veg Sci* 18:571–582. <https://doi.org/10.1111/j.1654-1103.2007.tb02571.x>
- Ghirardini MP, Carli M, Del Vecchio N, Rovati A, Cova O, Valigi F, Agnetti G, Macconi M, Adamo D, Traina M, Laudini F, Marcheselli I, Caruso N, Gedda T, Donati F, Marzadro A, Russi P, Spaggiari C, Bianco M, Binda R, Barattieri E, Tognacci A, Girardo M, Vaschetti L, Caprino P, Sesti E, Andreozzi G, Coletto E, Belzer G, Pieroni A (2007) The importance of a taste. A comparative study on wild food plant consumption in twenty-one local communities in Italy. *J Ethnobiol Ethnomed* 3:22. <https://doi.org/10.1186/1746-4269-3-22>
- Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A (2021) The world checklist of vascular plants, a continuously updated resource for exploring global plant diversity. *Sci Data* 8:215. <https://doi.org/10.1038/s41597-021-00997-6>
- Guarrera PM, Savo V (2016) Wild food plants used in traditional vegetable mixtures in Italy. *J Ethnopharmacol* 185:202–234. <https://doi.org/10.1016/j.jep.2016.02.050>
- Harris A (2023) An analysis of the 1977 za'atar regulation in Israel and Palestine through a lens of political ecology. Master's thesis, University of Oregon, Eugene
- Heywood V, Casas A, Ford-Lloyd B, Kell S, Maxted N (2007) Conservation and sustainable use of crop wild relatives. *Agric Ecosyst Environ* 121:245–255. <https://doi.org/10.1016/j.agee.2006.12.014>
- ISE (2006) International Society of Ethnobiology Code of Ethics (with 2008 additions). International Society of Ethnobiology. <https://www.ethnobiology.net/what-we-do/core-programs/ise-ethics-program/code-of-ethics/>. Accessed 22 August 2026
- IUCN (2026) The IUCN Red List of Threatened Species. Version 2026–1. <https://www.iucnredlist.org>. Accessed 22 August 2026
- Joly A, Bonnet P, Goëau H, Barbe J, Selmi S, Champ J, Dufour-Kowalski S, Affouard A, Carré J, Molino JF, Boujema N, Barthélémy D (2016) A look inside the PI@ntNet experience. *Multim Syst* 22:751–766. <https://doi.org/10.1007/s00530-015-0462-9>
- Landor-Yamagata JL, Kowarik I, Fischer LK (2018) Urban foraging in Berlin: people, plants and practices within the metropolitan green infrastructure. *Sustainability* 10:1873. <https://doi.org/10.3390/su10061873>
- Landucci F, Panella L, Lucarini D, Gigante D, Donnini D, Kell S, Maxted N, Venanzoni R, Negri V (2014) A prioritized inventory of crop wild relatives and wild harvested plants of Italy. *Crop Sci* 54:1628–1644. <https://doi.org/10.2135/cropsci2013.05.0355>
- Lenoir J, Gégout JC, Marquet PA, de Ruffray P, Brisse H (2008) A significant upward shift in plant species optimum elevation during the 20th century. *Science* 320:1768–1771. <https://doi.org/10.1126/science.1156831>
- Liguria R (2009) Legge regionale 10 luglio 2009, n. 28. Disposizioni in materia di tutela e valorizzazione della biodiversità. Bollettino Ufficiale della Regione Liguria n. 13 of 15 July 2009
- Łuczaj Ł, Pieroni A, Tardío J, Pardo-de-Santayana M, Sõukand R, Svanberg I, Kalle R (2012) Wild food plant use in 21st century Europe: the disappearance of old traditions and the search for new cuisines involving wild edibles. *Acta Soc Bot Pol* 81:359–370. <https://doi.org/10.5586/asbp.2012.031>
- Mattalia G, Quave CL, Pieroni A (2013) Traditional uses of wild food and medicinal plants among Brigasc, Kyé, and Provençal communities on the Western Italian Alps. *Genet Resour Crop Evol* 60:587–603. <https://doi.org/10.1007/s10722-012-9859-x>
- Maxted N, Ford-Lloyd BV, Jury S, Kell S, Scholten M (2006) Towards a definition of a crop wild relative. *Biodivers Conserv* 15:2673–2685. <https://doi.org/10.1007/s10531-005-5409-6>
- Maxted N, Kell S, Ford-Lloyd B, Dulloo E, Toledo Á (2012) Toward the systematic conservation of global crop wild relative diversity. *Crop Sci* 52:774–785. <https://doi.org/10.2135/cropsci2011.08.0415>
- McLain RJ, Hurley PT, Emery MR, Poe MR (2014) Gathering “wild” food in the city: rethinking the role of foraging in urban ecosystem planning and management. *Local Environ* 19:220–240. <https://doi.org/10.1080/13549839.2013.841659>
- Menendez-Baceta G, Aceituno-Mata L, Tardío J, Reyes-García V, Pardo-de-Santayana M (2012) Wild edible plants traditionally gathered in Gorbalea (Biscay, Basque Country). *Genet Resour Crop Evol* 59:1329–1347. <https://doi.org/10.1007/s10722-011-9760-z>
- Mina G, Scariot V, Peira G, Lombardi G (2023) Foraging practices and sustainable management of wild food resources in Europe: a systematic review. *Land* 12:1299. <https://doi.org/10.3390/land12071299>
- Niedrist G, Tasser E, Lüth C, Dalla Via J, Tappeiner U (2009) Plant diversity declines with recent land use changes in European Alps. *Plant Ecol* 202:195–210. <https://doi.org/10.1007/s11258-008-9487-x>
- Orsenigo S, Fenu G, Gargano D, Montagnani C, Abeli T, Alesandrini A, Bacchetta G, Bartolucci F, Carta A, Castello M, Cogoni D, Conti F, Domina G, Foggi B, Gennai M, Gigante D, Iberite M, Peruzzi L, Pinna MS, Prosser F, Santangelo A, Selvaggi A, Stinca A, Villani M, Wagen-sommer RP, Tartaglini N, Duprè E, Blasi C, Rossi G (2021) Red list of threatened vascular plants in Italy. *Plant Biosyst* 155:310–335. <https://doi.org/10.1080/11263504.2020.1739165>
- Parolo G, Rossi G (2008) Upward migration of vascular plants following a climate warming trend in the Alps. *Basic Appl Ecol* 9:100–107. <https://doi.org/10.1016/j.baee.2007.01.005>
- Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M, Benito Alonso JL, Coldea G, Dick J, Erschbamer B, Fernández Calzado R, Ghosn D, Holten JI, Kanka

- R, Kazakis G, Kollár J, Larsson P, Moiseev P, Moiseev D, Molau U, Grabherr G (2012) Recent plant diversity changes on Europe's mountain summits. *Science* 336:353–355. <https://doi.org/10.1126/science.1219033>
- Perrino EV, Perrino P (2020) Crop wild relatives: know how past and present to improve future research, conservation and utilization strategies, especially in Italy: a review. *Genet Resour Crop Evol* 67:1067–1105. <https://doi.org/10.1007/s10722-020-00930-7>
- Piemonte R (1982) Legge regionale 2 novembre 1982, n. 32. Norme per la conservazione del patrimonio naturale e dell'assetto ambientale. Bollettino Ufficiale della Regione Piemonte n. 45 of 10 November 1982
- Pieroni A, Giusti ME (2009) Alpine ethnobotany in Italy: traditional knowledge of gastronomic and medicinal plants among the Occitans of the upper Varaita valley, Piedmont. *J Ethnobiol Ethnomed* 5:32. <https://doi.org/10.1186/1746-4269-5-32>
- Pieroni A, Alrhoun M, Sulaiman N (2025) Plural and commoning? forecasting four scenarios for ethnobiology and ethnomedicine by 2035. *J Ethnobiol Ethnomed* 21:46. <https://doi.org/10.1186/s13002-025-00804-1>
- Pignatti S, Guarino R, La Rosa M (2017–2019) *Flora d'Italia*, 2nd edn, vols. 1–4. Edagricole – New Business Media, Bologna, Italy
- Poe MR, McLain RJ, Emery M, Hurley PT (2013) Urban forest justice and the rights to wild foods, medicines, and materials in the city. *Hum Ecol* 41:409–422. <https://doi.org/10.1007/s10745-013-9572-1>
- Reyes-García V, Menendez-Baceta G, Aceituno-Mata L, Acosta-Naranjo R, Calvet-Mir L, Domínguez P, Garnatje T, Gómez-Baggethun E, Molina-Bustamante M, Molina M, Rodríguez-Franco R, Serrasolses G, Vallès J, Pardo-Santayana M (2015) From famine foods to delicatessen: interpreting trends in the use of wild edible plants through cultural ecosystem services. *Ecol Econ* 120:303–311. <https://doi.org/10.1016/j.ecolecon.2015.11.003>
- Rumpf SB, Hülber K, Klöner G, Moser D, Schütz M, Wesely J, Willner W, Zimmermann NE, Dullinger S (2018) Range dynamics of mountain plants decrease with elevation. *Proc Natl Acad Sci USA* 115:1848–1853. <https://doi.org/10.1073/pnas.1713936115>
- Sansanelli S, Tassoni A (2014) Wild food plants traditionally consumed in the area of Bologna (Emilia Romagna region, Italy). *J Ethnobiol Ethnomed* 10:69. <https://doi.org/10.1186/1746-4269-10-69>
- Schunko C, Grasser S, Vogl CR (2015) Explaining the resurgent popularity of the wild: motivations for wild plant gathering in the Biosphere Reserve Grosses Walsertal, Austria. *J Ethnobiol Ethnomed* 11:55. <https://doi.org/10.1186/s13002-015-0032-4>
- Schunko C, Wild AS, Brandner A (2021) Exploring and limiting the ecological impacts of urban wild food foraging in Vienna. *Austria Urban for Urban Green* 62:127164. <https://doi.org/10.1016/j.ufug.2021.127164>
- Shackleton CM, Hurley PT, Dahlberg AC, Emery MR, Nagendra H (2017) Urban foraging: a ubiquitous human practice overlooked by urban planners, policy, and research. *Sustainability* 9:1884. <https://doi.org/10.3390/su9101884>
- Soga M, Gaston KJ (2016) Extinction of experience: the loss of human–nature interactions. *Front Ecol Environ* 14:94–101. <https://doi.org/10.1002/fee.1225>
- Steinbauer MJ, Grytnes JA, Jurasinski G, Kulonen A, Lenoir J, Pauli H, Rixen C, Winkler M, Bardy-Durchhalter M, Barni E, Bjorkman AD, Breiner FT, Burg S, Czortek P, Dawes MA, Delimat A, Dullinger S, Erschbamer B, Felde VA, Wipf S (2018) Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature* 556:231–234. <https://doi.org/10.1038/s41586-018-0005-6>
- Storkey J, Meyer S, Still KS, Leuschner C (2012) The impact of agricultural intensification and land-use change on the European arable flora. *Proc R Soc B* 279:1421–1429. <https://doi.org/10.1098/rspb.2011.1686>
- Sulaiman N, Aziz MA, Stryamets N, Mattalia G, Zocchi DM, Ahmed HM, Manduzai AK, Shah AA, Faiz A, Sökand R, Polesny Z, Pieroni A (2023) The importance of becoming tamed: wild food plants as possible novel crops in selected food-insecure regions. *Horticulturae* 9:171. <https://doi.org/10.3390/horticulturae9020171>
- Sulaiman N, Zocchi DM, Bonafede S, Nanni C, Sökand R, Pieroni A (2024a) Going or returning to nature? Wild vegetable uses in the foraging-centered restaurants of Lombardy, Northern Italy. *Plants* 13:2151. <https://doi.org/10.3390/plants13152151>
- Sulaiman N, Zocchi DM, Borrello MT, Mattalia G, Antoniazzi L, Berlinghof SE, Bewick A, Häfliger I, Schembs M, Torri L, Pieroni A (2024b) Traditional knowledge evolution over half of a century: local herbal resources and their changes in the upper Susa Valley of Northwest Italy. *Plants* 13:43. <https://doi.org/10.3390/plants13010043>
- Sulaiman N, Sökand R, Ullah I, Pieroni A (2026) Searching for a ghost?! the vain ethnobotany of foraging in three coastal Mediterranean areas. *J Ethnobiol Ethnomed* 22:15. <https://doi.org/10.1186/s13002-026-00853-0>
- Synk CM, Kim BF, Davis CA, Harding J, Rogers V, Hurley PT, Emery MR, Nachman KE (2017) Gathering Baltimore's bounty: characterizing behaviors, motivations, and barriers of foragers in an urban ecosystem. *Urban for Urban Green* 28:97–102. <https://doi.org/10.1016/j.ufug.2017.10.007>
- Tasser E, Walde J, Tappeiner U, Teutsch A, Noggler W (2007) Land-use changes and natural reforestation in the Eastern Central Alps. *Agric Ecosyst Environ* 118:115–129. <https://doi.org/10.1016/j.agee.2006.05.004>
- Ticktin T (2004) The ecological implications of harvesting non-timber forest products. *J Appl Ecol* 41:11–21. <https://doi.org/10.1111/j.1365-2664.2004.00859.x>
- Turner NJ, Łuczaj ŁJ, Migliorini P, Pieroni A, Dreon AL, Sacchetti LE, Paoletti MG (2011) Edible and tended wild plants, traditional ecological knowledge and agroecology. *Crit Rev Plant Sci* 30:198–225. <https://doi.org/10.1080/07352689.2011.554492>
- Vincent H, Wiersema J, Kell S, Fielder H, Dobbie S, Castañeda-Álvarez NP, Guarino L, Eastwood R, León B, Maxted N (2013) A prioritized crop wild relative inventory to help underpin global food security. *Biol Conserv* 167:265–275. <https://doi.org/10.1016/j.biocon.2013.08.011>

- Vitalini S, Iriti M, Puricelli C, Ciuchi D, Segale A, Fico G (2013) Traditional knowledge on medicinal and food plants used in Val San Giacomo (Sondrio, Italy): an alpine ethnobotanical study. *J Ethnopharmacol* 145:517–529. <https://doi.org/10.1016/j.jep.2012.11.024>
- von Wettberg E, Davis TM, Smýkal P (2020) Editorial: wild plants as source of new crops. *Front Plant Sci* 11:591554. <https://doi.org/10.3389/fpls.2020.591554>
- Zocchi DM, Fontefrancesco MF, Corvo P, Pieroni A (2021) Recognising, safeguarding, and promoting food heritage: challenges and prospects for the future of sustainable food systems. *Sustainability* 13:9510. <https://doi.org/10.3390/su13179510>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.